

**Impacts of Atmospheric Anthropogenic Nitrogen on the Open Ocean**R. A. Duce, *et al.**Science* **320**, 893 (2008);

DOI: 10.1126/science.1150369

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of March 1, 2012):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/320/5878/893.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2008/05/15/320.5878.893.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/320/5878/893.full.html#related>

This article has been **cited by** 68 article(s) on the ISI Web of Science

This article has been **cited by** 9 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/320/5878/893.full.html#related-urls>

This article appears in the following **subject collections**:

Oceanography

<http://www.sciencemag.org/cgi/collection/oceans>

Impacts of Atmospheric Anthropogenic Nitrogen on the Open Ocean

R. A. Duce,^{1*} J. LaRoche,² K. Altieri,³ K. R. Arrigo,⁴ A. R. Baker,⁵ D. G. Capone,⁶ S. Cornell,⁷ F. Dentener,⁸ J. Galloway,⁹ R. S. Ganeshram,¹⁰ R. J. Geider,¹¹ T. Jickells,⁵ M. M. Kuypers,¹² R. Langlois,² P. S. Liss,⁵ S. M. Liu,¹³ J. J. Middelburg,¹⁴ C. M. Moore,¹¹ S. Nickovic,¹⁵ A. Oschlies,² T. Pedersen,¹⁶ J. Prospero,¹⁷ R. Schlitzer,¹⁸ S. Seitzinger,³ L. L. Sorensen,¹⁹ M. Uematsu,²⁰ O. Ulloa,²¹ M. Voss,²² B. Ward,²³ L. Zamora¹⁷

Increasing quantities of atmospheric anthropogenic fixed nitrogen entering the open ocean could account for up to about a third of the ocean's external (nonrecycled) nitrogen supply and up to ~3% of the annual new marine biological production, ~0.3 petagram of carbon per year. This input could account for the production of up to ~1.6 teragrams of nitrous oxide (N₂O) per year. Although ~10% of the ocean's drawdown of atmospheric anthropogenic carbon dioxide may result from this atmospheric nitrogen fertilization, leading to a decrease in radiative forcing, up to about two-thirds of this amount may be offset by the increase in N₂O emissions. The effects of increasing atmospheric nitrogen deposition are expected to continue to grow in the future.

Nitrogen is an essential nutrient in terrestrial and marine ecosystems. Most nitrogen in the atmosphere and ocean is present as N₂ and is available only to diazotrophs, a restricted group of microorganisms that can fix

N₂. Most organisms can only assimilate forms of reactive nitrogen (fixed nitrogen, N_r), including oxidized and reduced inorganic and organic forms. The availability of N_r limits primary production, the conversion of inorganic carbon to organic carbon (*I*), in much of the ocean. Reactive nitrogen enters the ocean via rivers, N₂ fixation, and atmospheric deposition. It is removed via N₂ formation by denitrification and anaerobic ammonium oxidation (anammox), nitrous oxide (N₂O) and ammonia emissions, and burial of organic matter in sediments. Human activities have severely altered many coastal ecosystems by increasing the input of anthropogenic nitrogen through rivers and groundwater, direct discharges from wastewater treatment, atmospheric deposition, and so forth, resulting in increasing eutrophication. Human activities have also added large quantities of atmospheric N_r to central ocean regions.

Riverine input of N_r to the oceans is estimated as 50 to 80 Tg N year⁻¹ (2–4). However, much is either lost to the atmosphere after N₂ conversion or buried in coastal sediments, never reaching oceanic regions (5). We assume that riverine N_r has a negligible impact on the open ocean nitrogen inventory, and we do not consider it further. Estimates of global ocean N₂ fixation range from 60 to 200 Tg N year⁻¹ (2, 6–8). Although impacts of the amplified nitrogen inputs to terrestrial systems are being continuously evaluated (3, 9), here we show that atmospheric transport and deposition is an increasingly important pathway for N_r entering the open ocean, often poorly represented in analyses of open ocean anthropogenic impacts (10–16). Atmospheric N_r input is rapidly approaching global oceanic estimates for N₂ fixation and is predicted to increase further due to emissions from combustion of fossil fuels and production and use of fertilizers. Our objective is to highlight the growing im-

portance of anthropogenic atmospheric N_r (AAN) deposition to the oceans and evaluate its impact on oceanic productivity and biogeochemistry.

Atmospheric Emission and Deposition of Nitrogen Species

Atmospheric emissions of N_r are primarily oxidized nitrogen species, NO_x (NO + NO₂) and NH₃. Recent studies suggest that atmospheric water-soluble organic nitrogen is far more abundant than conventionally thought, constituting ~30% of total N_r deposition (13, 17–20). Given the uncertain origins and complex composition of this material, the importance of direct emissions and secondary formation of organic nitrogen is unclear. However, measurements suggest that an important fraction is anthropogenic (13, 17). We therefore assume that in 1860, the relationship between organic and inorganic nitrogen deposition was the same as it is today and increase our 1860 estimate so that organic nitrogen represents 30% of total N_r deposition. The uncertainties associated with this assumption emphasize the need for further research on atmospheric organic nitrogen.

Estimated total N_r and AAN emissions in 1860, 2000, and 2030 (Table 1) show that anthropogenic emissions have significantly increased since the mid-1800s and future increases are expected (21). Over the next 20 to 25 years, the proportion of NH₃ emissions will likely increase due to enhanced atmospheric emission controls predicted to be more effective for NO_x than NH₃ (Table 1) (21). An important fraction of atmospheric N_r emissions is deposited on the ocean (Table 1). In 1860, this amounted to ~20 Tg N year⁻¹, of which ~29% was anthropogenic. By 2000, the total N_r deposition to the ocean had more than tripled to ~67 Tg N year⁻¹, with ~80% being anthropogenic. This is greater than the 39 Tg N year⁻¹ reported by (14), in part because our estimate includes water-soluble organic nitrogen. Estimates of anthropogenic emissions for 2030 indicate a ~4-fold increase in total atmospheric N_r deposition to the ocean and an ~11-fold increase in AAN deposition compared with 1860 (22).

The spatial distribution of atmospheric deposition has also changed greatly (Fig. 1, A and B). Deposition to most of the ocean was <50 mg N m⁻² year⁻¹ in 1860, with very few areas >200 mg N m⁻² year⁻¹. Most oceanic deposition was from natural sources; anthropogenic sources impacted only a few coastal regions. By 2000, deposition over large ocean areas exceeded 200 mg N m⁻² year⁻¹, reaching >700 mg N m⁻² year⁻¹ in many areas. Intense deposition plumes extend far downwind of major population centers in Asia, India, North and South America, around Europe, and west of Africa (Fig. 1B). A direct comparison of deposition in 1860 and 2000 shows almost all ocean surface areas now being affected by AAN deposition (Fig. 1, A and B). Predictions for 2030 (fig. S1) indicate similar patterns, but with

¹Departments of Oceanography and Atmospheric Sciences, Texas A&M University, College Station, TX 77843, USA. ²Leibniz-Institut fuer Meereswissenschaften, 24105 Kiel, Germany. ³Institute of Marine and Coastal Sciences, Rutgers University, Rutgers/NOAA CMER Program, New Brunswick, NJ 08901, USA. ⁴Department of Environmental Earth System Science, Stanford University, Stanford, CA 94305, USA. ⁵School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK. ⁶Department of Biological Sciences, University of Southern California, Los Angeles, CA 90089, USA. ⁷QUEST—Earth Sciences, University of Bristol, Bristol BS8 1RJ, UK. ⁸European Commission, Joint Research Centre, Institute for Environment and Sustainability, TP290, I-21020, Ispra (Va), Italy. ⁹Department of Environmental Sciences, University of Virginia, Charlottesville, VA 22904, USA. ¹⁰John Murray Laboratories, The King's Buildings, Edinburgh EH9 3JW, UK. ¹¹Department of Biological Sciences, University of Essex, Colchester CO4 3SQ, UK, and National Oceanography Centre, University of Southampton, Southampton SO14 3ZH, UK. ¹²Max Planck Institute for Marine Microbiology, Celsiusstrasse 1, D-28359 Bremen, Germany. ¹³Key Laboratory of Marine Chemistry Theory and Technology Ministry of Education, College of Chemistry and Chemical Engineering, Ocean University of China, Qingdao 266100, Peoples Republic of China. ¹⁴Netherlands Institute of Ecology, Korringaweg 7, 4401 NT Yerseke, Netherlands. ¹⁵Atmospheric Research and Environment Programme, World Meteorological Organization, BP2300, 1211 Geneva 2, Switzerland. ¹⁶University of Victoria, Post Office Box 3055 STN CSC, Victoria, BC V8W 3P6, Canada. ¹⁷Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, Miami, FL 33149, USA. ¹⁸Alfred Wegener Institute for Polar and Marine Research, 27568 Bremerhaven, Germany. ¹⁹National Environmental Research Institute, Aarhus University, Denmark. ²⁰Ocean Research Institute, University of Tokyo, Tokyo 164-8639, Japan. ²¹Departamento de Oceanografía, Centro de Investigación Oceanográfica, COPAS, and Nucleo Milenio EMBA, Universidad de Concepción, Casilla 160-C, Concepción, Chile. ²²Leibniz Institute for Baltic Sea Research, Warnemünde, 18119 Rostock, Germany. ²³Department of Geosciences, Princeton University, Princeton, NJ 08544, USA.

*To whom correspondence should be addressed. E-mail: rduce@ocean.tamu.edu

increased deposition further into open ocean regions (21, 22). The ratio of 2030-to-2000 deposition rates (Fig. 1C) shows up to a factor of 2 increase in Southeast Asia, the Bay of Bengal, and the Arabian Sea; up to a 50% increase off western Africa; and up to 30% across essentially all the mid-latitude North Atlantic and North Pacific. As Galloway *et al.* (9) conclude, controlling NO_x emissions using maximum feasible reductions could substantially decrease future emissions, so

the increases we predict on deposition rates (Fig. 1C) may represent upper limits.

Impact on New Primary Production and the Biological Pump

Present global open ocean primary production is estimated at $\sim 50 \text{ Pg C year}^{-1}$ (23), equivalent to $\sim 8800 \text{ Tg N year}^{-1}$, assuming Redfield stoichiometry (Table 2). Because $\sim 78\%$ of this production is driven by regeneration of N_r within surface

waters (24) (*a* in Fig. 2), it is more relevant to evaluate the impact of AAN deposition on oceanic productivity and biogeochemistry by comparing AAN with global new production, estimated at $\sim 11 \text{ Pg C year}^{-1}$ (24–26). New production (*b* in Fig. 2 and Table 2) is dominated by nitrate regenerated at depth from sinking organic matter and subsequently returned to the euphotic zone via physical transport (*b'* in Fig. 2) (27). Over sufficiently large space and time scales




Fig. 1. (A) Total atmospheric reactive nitrogen (N_r) deposition in 1860 in $\text{mg m}^{-2} \text{ year}^{-1}$ [NH_x and NO_y are derived from (3), with the addition of 30% of the total nitrogen as organic nitrogen]. Total atmospheric N_r deposition in 1860 was $\sim 20 \text{ Tg N year}^{-1}$, AAN was $\sim 5.7 \text{ Tg N year}^{-1}$. (B) Total atmospheric reactive nitrogen (N_r) deposition in 2000 in $\text{mg m}^{-2} \text{ year}^{-1}$ [derived from (21) with the addition of 30% of the total nitrogen as organic nitrogen]. Total atmospheric N_r deposition in 2000 was $\sim 67 \text{ Tg N year}^{-1}$, AAN was $\sim 54 \text{ Tg N year}^{-1}$. (C) Ratio of the projected flux of N_r to the ocean in 2030 to

that in 2000. (D) Nitrate concentrations (μM) in the surface (0 to 1 m) waters of the ocean (43). (E) Similar to (B), but with regions where surface nitrate $> 4 \mu\text{M}$ has been masked out. Total atmospheric N_r deposition in 2000 to the nonmasked areas was $\sim 51 \text{ Tg N year}^{-1}$, AAN was $\sim 41 \text{ Tg N year}^{-1}$. (F) Ratio of total N_r deposition to dissolved inorganic nitrogen (DIN) supply into the upper 130 m as diagnosed from a model fitted to oceanic tracer observations (44). To reduce noise, computation of the ratio has been limited to areas with DIN supply exceeding $0.05 \text{ mol m}^{-2} \text{ year}^{-1}$.

(1 to ~1000 years), nitrate-driven new production is balanced by the biologically mediated export of particulate and dissolved organic matter from the surface layer (b'' in Fig. 2). On a similar time scale, this component of new production is almost neutral in terms of carbon assimilation (28) because degradation processes release N_r and CO_2 in stoichiometric amounts equivalent to the initial elemental composition of the organic matter. In the absence of denitrification and other fixed nitrogen losses in the ocean interior, nitrate-based new production can be considered a closed loop within which the biologically mediated carbon export (b'') is balanced by a return flux of dissolved inorganic carbon (b'), resulting in near-zero net air-sea CO_2 exchange.

Only external (to the ocean) sources of N_r that reach the surface mixed layer can affect the steady-state balance of the biologically mediated flux of CO_2 across the air-sea interface. The two known open ocean sources of external N_r are biological N_2 fixation (c in Fig. 2) and atmospheric deposition (d). Together these contribute a net oceanic input of N_r that can support “completely new production” and hence influence global oceanic N_r and the net atmosphere-to-ocean exchange of CO_2 , assuming an adequate supply of other nutrients (P, Fe). Although N_2 fixation must have dominated the flux of external new nitrogen in the preindustrial world, atmospheric N_r deposition is now approaching N_2 fixation as a result of the

dramatic increase in the anthropogenic component (Table 2).

Can this atmospheric N_r deposition be rapidly assimilated into primary production? It will impact the biogeochemistry of oceanic areas that are either perennially or seasonally depleted in surface nitrate, but will have little effect in high-nutrient, low-chlorophyll (HNLC) regions where the concentration of surface nitrate is always high. Comparing surface nitrate concentrations (Fig. 1D) and total N_r deposition (Fig. 1B) shows the relatively small overlap between high N_r deposition and significant surface nitrate concentrations. In regions where surface nitrate is seasonally depleted (i.e., where productivity is nitrogen limited), atmospheric deposition would likely be assimilated during the year. Although N_r generally is seasonally exhausted in regions where mean annual nitrate is $<7 \mu M$, a more conservative value of $<4 \mu M$ is used to calculate the distribution of the atmospheric N_r deposition in present-day nitrogen-depleted waters (Fig. 1E). The calculated global N_r deposition to regions with mean nitrate $<4 \mu M$ is $\sim 51 \text{ Tg N year}^{-1}$, or $\sim 76\%$ of the total atmospheric N_r deposited in the ocean, compared to $\sim 56 \text{ Tg N year}^{-1}$ ($\sim 84\%$ of total deposition) if $<7 \mu M$ is used as a threshold. Corresponding values for AAN are ~ 41 and $\sim 45 \text{ Tg N year}^{-1}$. Using the areas delineated by the $<4 \mu M$ and $<7 \mu M$ nitrate concentrations above, we calculate that ~ 67 to 75% of oceanic

surface waters are potentially seasonally nitrogen limited, although some of these areas may not be exclusively nitrogen limited but rather colimited (1). It has recently been assumed that only 40% of the ocean is nitrogen limited (14), although this estimate did not allow for N/P colimitation such as seen in the North Atlantic and other areas designated P-limited in (14). These are likely underestimates because much of the N_r is deposited upstream of N_r -depleted regions (e.g., HNLC Southern Ocean) and will eventually be advected into thermocline waters of nitrogen-limited regions of the Southern Hemisphere and North Atlantic and thus are important to future (decades to centuries) productivity and biogeochemistry (29).

The total atmospheric deposition plus N_2 -fixation flux to the ocean is $\sim 167 \text{ Tg N year}^{-1}$ (Table 2). Assuming complete assimilation, these external N_r sources can support a maximum biologically mediated flux of $\sim 1.0 \text{ Pg C year}^{-1}$, of which $\sim 0.4 \text{ Pg C year}^{-1}$ is from atmospheric deposition. Deposition of AAN alone could support up to $\sim 0.3 \text{ Pg C year}^{-1}$, or $\sim 3\%$ of all new production, including that from nutrients upwelled from deep waters, and $\sim 32\%$ of the productivity derived from external N_r supply (Table 2). In 1860, AAN supported a biologically mediated carbon flux of only $\sim 0.03 \text{ Pg C year}^{-1}$, so from 1860 to the present the potential impact of AAN on net primary productivity has increased ~ 10 -fold. An earlier lower estimate ($0.16 \text{ Pg C year}^{-1}$) of new (export) production generated by AAN deposition (14) assumed a different nitrogen-limited area, lower atmospheric fluxes, and the assumption that N enhancement will result in the replacement of diazotrophs by other phytoplankton.

Increased new production due to AAN fertilization coincides with the anthropogenic perturbation of the global carbon cycle and penetration of anthropogenic carbon in the ocean. The current anthropogenic CO_2 uptake by the ocean is $\sim 2.2 \pm 0.5 \text{ Pg C year}^{-1}$ (30), primarily attributed to physical-chemical processes (the “solubility pump”). Assuming that new production draws down atmospheric CO_2 according to Redfieldian stoichiometry, up to $\sim 10\%$ of the present anthropogenic carbon uptake could be attributed to anthropogenic nitrogen fertilization. This potentially significant enhancement of the oceanic uptake of anthropogenic carbon indicates the need to incorporate this factor in future Earth system assessments and models, as has already been done for terrestrial ecosystems (31). This estimate may be lower if the dissolved organic carbon or particulate organic carbon produced is regenerated at shallow depths (32). The efficiency and longevity of this anthropogenic nitrogen fertilization effect depend on temporal uncoupling of the new N_r inputs (N_2 fixation and atmospheric deposition) from N_r removal (e.g., denitrification/anammox and burial). Assuming that all other essential nutrients are in adequate supply, it will be operational as long as the




Fig. 2. Schematic of the processes supplying nutrients for surface primary production. See text for detailed description.

increase in new N_r (and associated additional CO_2 uptake) is not balanced by increased regeneration of N_2 and CO_2 and release at the ocean-air interface. Eventually, if AAN deposition levels off, the ocean may reach a new steady state with respect to nitrogen gains and losses that is neutral with respect to CO_2 uptake over time scales similar to the oceanic N residence time (~1000 years).

The future impact of AAN on productivity must be evaluated in the context of predicted changes in productivity caused by other variables. For instance, elevated concentrations of atmospheric CO_2 may have resulted in excess carbon consumption and export because of shifting C:N stoichiometry (33), and it is unclear whether projected AAN and high CO_2 concentrations have synergy or compensate. El Niño–Southern Oscillation (ENSO)–induced higher

water temperatures and the associated increased stratification in low-latitude oceans may have reduced productivity by 60% in some regions (34). Thus, in a warmer climate, decreases in productivity due to restricted injection of nutrient-rich deep water would only accentuate the importance of AAN contributions to new production in low-latitude oligotrophic oceanic areas where AAN already has a strong effect. Assuming that all N_r deposition is assimilated into primary production, this N_r -driven new production could contribute as much as 20% of the total new (or export) production in such regions where upwelling is limited, e.g., the North Atlantic gyre (Fig. 1F). The contribution of N_r deposition to new production is higher in the Atlantic than the Pacific and can reach magnitudes comparable to export production along some continental areas.

Table 1. Atmospheric nitrogen emissions and deposition to the ocean. Assumed uncertainties—emissions: 1860: $\pm 50\%$; 2000: $NO_x \pm 30\%$, $NH_3 \pm 50\%$; 2030: see text and (20). Deposition: 1860: $\pm 50\%$; 2000: $NO_y/NH_x \pm 40\%$, organic N $\pm 50\%$; 2030: see text and (20).

	1860* (Tg N year ⁻¹)	2000† (Tg N year ⁻¹)	2030‡ (Tg N year ⁻¹)
<i>Emission to the atmosphere</i>			
Total NO_x	13 (7–20)	52 (36–68)	54‡
Anthropogenic NO_x	2.6 (1.3–4)	38 (27–49)	43
Total NH_3	21 (11–32)	64 (32–96)	78‡
Anthropogenic NH_3	7.4 (3.7–11)	53 (27–80)	70
Total atmospheric N emissions	34 (18–52)	116 (68–164)	132
Total anthropogenic N_r (AAN)	10 (5–15)	91 (54–129)	113
<i>Deposition to the ocean</i>			
Total NO_y	6.2 (3.1–9.3)	23 (14–32)	25
Anthropogenic NO_y	1.2 (0.6–1.8)	17 (10–24)	18
Total NH_x	8 (4–12)	24 (14–34)	29
Anthropogenic NH_x	2.4 (1.2–3.6)	21 (13–29)	25
Total organic N_r	6.1 (3.0–9.1)	20 (10–30)	23
Anthropogenic organic N_r	2.1 (1.0–3.1)	16 (8–24)	19
Total N_r deposition	20 (10–30)	67 (38–96)	77
Total anthropogenic N_r (AAN)	5.7 (2.8–8.5)	54 (31–77)	62

*From (3). †Derived from (21); see text and (26). ‡ NO_x and NH_3 based on ~80% and ~90% anthropogenic, respectively [from (3)].

Table 2. Atmospheric nitrogen deposition to the ocean in 2000 and its impact on productivity. Global-scale estimates of total primary production (23); new production (24–26); N_2 fixation (2, 6–8). Most letters in italics refer to flux pathways in Fig. 2.

	Global ocean nitrogen (Tg N year ⁻¹)	Resultant global ocean productivity (Pg C year ⁻¹)
Total primary production (<i>a+b+c+d</i>)	~8800 (7000–10,500)	~50 (40–60)
New production (NP) (<i>b</i>)	~1900 (1400–2600)	~11 (8–15)
Marine N_2 fixation (<i>c</i>)	~100 (60–200)	~0.57 (0.3–1.1)
Total net N_r deposition (<i>d</i>) ($NO_y+NH_x+Org. N_r$)	~67 (38–96)	~0.38 (0.22–0.55)
Total external nitrogen supply (<i>c+d</i>)	~167 (98–296)	~0.95 (0.56–1.7)
Anthropogenic N_r deposition (AAN) (<i>e</i>)	~54 (31–77)	~0.31 (0.18–0.44)
Marine N_2 fixation as % NP N_r	= <i>c/b</i>	~5.3% (2.3–14.3%)
Total N_r deposition as % NP N_r	= <i>d/b</i>	~3.5% (1.5–6.9%)
AAN as % NP N_r	= <i>e/b</i>	~2.8% (1.2–5.5%)
Total N_r deposition as % external N supply	= <i>d/(c+d)</i>	~40% (13–98%)
AAN as % external N supply	= <i>e/(c+d)</i>	~32% (10–79%)

On the basis of future scenarios for anthropogenic emissions, AAN contribution to primary production could approach current estimates of global N_2 fixation by 2030. Fertilization of the surface layer by atmospheric deposition, primarily AAN, could even lead to a decrease in N_2 fixation due to biological competition (14). However, atmospheric N_r deposition has a very small effect on the surface seawater ambient N_r concentrations, too little to inhibit nitrogenase activity directly [e.g., we estimate that an extremely rare and large atmospheric deposition event distributed over a 25-m mixed-layer depth could increase the N_r concentration by only ~45 nM (35), which is too small to suppress N_2 fixation (36)]. Atmospheric N_r deposition more likely represents a long-term low-level fertilization of the ocean that has consequences for the natural biogeochemical cycles of nitrogen and carbon and their ongoing anthropogenic perturbations. Biological evidence suggests that phytoplankton communities in oceanic gyres are presently nitrogen limited (1). Atmospheric N_r deposition, in the absence of significant atmospheric deposition of phosphorus, may exacerbate phosphorus limitation of N_2 fixation. The long-term effect of AAN deposition on N_2 fixation depends on whether P or Fe limits N_2 fixation and on the supply ratio of bioavailable N:P:Fe derived from atmospheric deposition (37). Atmospheric deposition of phosphorus is much less perturbed by human activity than N_r (13, 37). Hence, the overall impact of atmospheric deposition is likely to be a shift in the N/P balance of surface waters. Some marine diazotrophs can exploit dissolved organic phosphorus pools and may obtain an adequate P supply by degrading compounds such as phosphonates (38).

Changes in species composition and productivity can lead to changes in the export of nitrogen and carbon to deep ocean water, resulting in a shift of deep ocean N/P ratios away from Redfield stoichiometry, which could then influence the chemistry of upwelled waters remote from the loci of atmospheric depositions. Remineralization of this extra organic carbon flux in deep waters may reduce the deepwater O_2 concentration, and the resultant microbial N_2 production will act to restore the N/P ratio toward the Redfield value, as suggested to have happened in the past (39). (See Supporting Online Material, including fig. S2).

Impact on N_2O Emissions from the Ocean

Another important issue is whether increasing atmospheric N_r inputs to the ocean can alter marine emissions of nitrous oxide (N_2O), a major greenhouse gas. Estimates of global sea-to-air N_2O fluxes vary considerably. Two recent estimates are the Intergovernmental Panel on Climate Change (IPCC) assessment (30) (3.8 Tg N year⁻¹ as N_2O) and the calculation by Bange of the mean from data in (40) (6.2 Tg N year⁻¹). Using the mean (5.0 Tg N year⁻¹) and the range of these two estimates, and assuming that the nitrogen in

this “recent” N_2O flux originally entered the oceans from N_2 fixation ($100 \text{ Tg N year}^{-1}$) and atmospheric deposition ($67 \text{ Tg N year}^{-1}$), then the emission of $5.0 \text{ Tg N year}^{-1}$ as N_2O results from nitrification and denitrification of part of this $167 \text{ Tg N year}^{-1}$ entering the surface ocean. This assumes that N_2O production in the near-surface ocean is at steady state and there are no significant time lags between atmospheric input and N_2O formation. Normalizing the N_2O flux to the atmosphere by the “completely new” nitrogen input (5.0:167) can then be used to estimate that AAN deposition has resulted in the production of up to $\sim 1.6 \text{ Tg N}_2\text{O-N year}^{-1}$, or about a third of total oceanic N_2O emissions. This approach suggests that in 1860, only $\sim 0.2 \text{ Tg N year}^{-1}$ ($\sim 5\%$) of the sea-to-air flux of N_2O was driven by atmospheric anthropogenic inputs, assuming simplistically that N_2O production is linearly related to N supply. [We use linear scaling due to the lack of experimental and modeling studies that address the spatial and nonlinear response of N_2O emissions to N deposition, although important regional variations are likely (41).] This suggests that from 1860 to the present, the increase in AAN has led to nearly an order of magnitude increase in anthropogenic N_2O emission from the oceans. Calculations and estimates of increases for 2030 are in table S1.

While oceanic AAN deposition may result in increased N_2O emissions, increasing radiative forcing, AAN also increases primary production (up to $\sim 0.3 \text{ Pg C year}^{-1}$ detailed above) and export production to the deep ocean, removing CO_2 from the atmosphere and therefore decreasing radiative forcing. With a Global Warming Potential of 298 for N_2O (42), the net balance suggests that about two-thirds of the decrease in radiative forcing from CO_2 uptake could be offset by the increase due to N_2O emissions. The uncertainty in our estimates is considerable; however, the estimates suggest the potential importance of AAN to N_2O emissions and therefore the need for future research in regions such as oceanic Oxygen Minimum Zones (OMZs), which, although small in area, are potentially important for N_2O emissions. The future role of OMZs will be influenced not only by AAN but also by climate and other global changes.

Conclusions

This analysis emphasizes the potential importance of the growing quantity of atmospheric reactive (fixed) nitrogen that enters the open ocean as a result of human activities and its impact on the present marine nitrogen cycle. Considering the increasing demand for energy and fertilizers, the emissions of AAN are expected to grow over the coming decades. Atmospheric deposition of anthropogenic nitrogen to the ocean may account for up to $\sim 3\%$ of the annual new oceanic primary productivity, but about a third of the primary productivity generated as a result of the external input of nitrogen to the ocean. The input of AAN is approaching that of

N_2 fixation as a source of marine reactive nitrogen. Although local AAN deposition seems unlikely to alter significantly local phytoplankton species composition, the phytoplankton community could be affected by the slow long-term fertilization of surface waters by AAN. Moreover, AAN inputs to the ocean have potentially important climatic implications. Up to about a tenth of the anthropogenic atmospheric carbon uptake by the ocean (as CO_2) may result from this fertilization. In addition, AAN inputs may stimulate N_2O emissions, with possibly about two-thirds of the decrease in radiative forcing from increased CO_2 uptake by the ocean being offset by the increase in radiative forcing from increased N_2O emissions.

There is clearly much we do not know about the extent and time scale of the impacts of AAN deposition on the oceans and the feedbacks to the climate system. The issues are complex and interactive, and they must be considered in climate scenarios. Areas of particular importance include understanding more fully the sources, chemical speciation, reactivity, and availability of atmospheric organic nitrogen; developing more realistic models of N_r deposition to the ocean, coupled with measuring N_r deposition over extended periods of time in open ocean regions; understanding the relationships between, and impacts of, the atmospheric deposition of bioavailable N, P, and Fe; and understanding the mechanisms and time scales involved in the oceanic response to N_r deposition, coupled with a new generation of Earth system models that take into account long-term low-level nitrogen fertilization of the ocean and evaluate the effect on N_2O emissions and the duration of the enhanced (anthropogenic) CO_2 uptake.

References and Notes

- M. M. Mills, C. Ridame, M. Davey, J. La Roche, R. J. Geider, *Nature* **429**, 292 (2004).
- N. Gruber, J. Sarmiento, in *The Sea: Biological-Physical Interactions*, A. R. Robinson, J. F. McCarthy, B. Rothschild, Eds. (Wiley, New York, 2002), vol. 12, pp. 337–399.
- J. N. Galloway *et al.*, *Biogeochemistry* **35**, 3 (1996).
- S. P. Seitzinger, J. A. Harrison, E. Dumont, A. H. W. Beusen, A. F. Bouwman, *Global Biogeochem. Cycles* **19**, GB4501 (2005).
- S. Seitzinger *et al.*, *Ecol. Appl.* **16**, 2064 (2006).
- C. Mahaffey, A. F. Michaels, D. G. Capone, *Am. J. Sci.* **305**, 546 (2005).
- J. K. Moore, S. C. Doney, K. Lindsay, N. Mahowald, A. F. Michaels, *Tellus* **58B**, 560 (2006).
- C. Deutsch, J. L. Sarmiento, D. M. Sigman, N. Gruber, J. P. Dunne, *Nature* **445**, 163 (2007).
- J. N. Galloway *et al.*, *Science* **320**, 889 (2008).
- R. A. Duce *et al.*, *Global Biogeochem. Cycles* **5**, 193 (1991).
- J. M. Prospero *et al.*, *Biogeochemistry* **35**, 27 (1996).
- C. M. Duarte *et al.*, *J. Geophys. Res.* **111**, G04006 (2006).
- T. Jickells, *Biogeosciences* **3**, 271 (2006).
- A. Krishnamurthy, J. K. Moore, C. S. Zender, C. Luo, *J. Geophys. Res.* **112**, G02019 (2007).
- H. W. Paerl, *Nature* **315**, 747 (1985).
- S. C. Doney *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **104**, 14580 (2007).
- S. E. Cornell, T. D. Jickells, J. N. Cape, A. P. Rowland, R. A. Duce, *Atmos. Environ.* **37**, 2173 (2003).
- K. A. Mace, R. A. Duce, N. W. Tindale, *J. Geophys. Res.* **108**, 4338 (2003).
- T. Nakamura, H. Ogawa, D. K. Maripi, M. Uematsu, *Atmos. Environ.* **40**, 7259 (2006).

- If the assumption that in 1860 the relationship between organic and inorganic N in deposition was the same as today, i.e., that organic nitrogen is $\sim 30\%$ of the total N_r , is in error, then it is likely that both the total and the anthropogenic nitrogen deposition in 1860 would have been less than indicated in Table 1.
- F. Dentener *et al.*, *Global Biogeochem. Cycles* **20**, GB4003 (2006).
- The deposition estimates for 2030 are based on the S2 simulation of NO_x and NH_3 emissions to the atmosphere presented in (21), which uses an IASA CLE 2030 current emission regulation scenario, termed “likely” in that paper. We estimate that the atmospheric emission and deposition values shown in Table 1 for 2030 have uncertainties of ~ 40 to 50% . Dentener *et al.* (21) also discuss results using the “optimistic” IASA Maximum Feasible Reduction (MFR) scenario and the “pessimistic” IPCC SRES A2 scenario. Depending on the regional development path of N, emissions, N, depositions may be lower by 10 to 70% (MFR) or higher by 30 to 200%.
- M. E. Carr *et al.*, *Deep-Sea Res. Part II* **53**, 741 (2006).
- E. A. Laws, P. G. Falkowski, W. O. Smith, H. Ducklow, J. J. McCarthy, *Global Biogeochem. Cycles* **14**, 1231 (2000).
- A. Oschlies, *Deep-Sea Res. Part II* **48**, 2173 (2001).
- H. W. Ducklow, *Rev. Geophys.* **33**, 1271 (1995).
- R. C. Dugdale, J. J. Goering, *Limnol. Oceanogr.* **12**, 196 (1967).
- W. S. Broecker, *Global Biogeochem. Cycles* **5**, 191 (1991).
- J. L. Sarmiento, N. Gruber, M. A. Brzezinski, J. P. Dunne, *Nature* **427**, 56 (2004).
- K. Denman *et al.*, in *Climate Change 2007: The Physical Science Basis*, S. Solomon *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 2007), pp. 544–547.
- F. Magnani *et al.*, *Nature* **447**, 849 (2007).
- A. Gnanadesikan, J. L. Sarmiento, R. D. Slater, *Global Biogeochem. Cycles* **17**, 1050 (2003).
- U. Riebesell *et al.*, *Nature* **450**, 545 (2007).
- M. J. Behrenfeld *et al.*, *Nature* **444**, 752 (2006).
- A. F. Michaels, D. A. Siegel, R. J. Johnson, A. H. Knap, J. N. Galloway, *Global Biogeochem. Cycles* **7**, 339 (1993).
- C. M. Holl, J. P. Montoya, *J. Phycol.* **41**, 1178 (2005).
- A. R. Baker, S. D. Kelly, K. F. Biswas, M. Witt, T. D. Jickells, *Geophys. Res. Lett.* **30**, 2296 (2003).
- S. T. Dyrhman *et al.*, *Nature* **439**, 68 (2006).
- R. S. Ganeshram, T. F. Pedersen, S. E. Calvert, G. W. McNeill, M. R. Fontugne, *Paleoceanography* **15**, 361 (2000).
- H. W. Bange, *Atmos. Environ.* **40**, 198 (2006).
- X. Jin, N. Gruber, *Geophys. Res. Lett.* **30**, GL18458 (2003).
- S. Solomon *et al.*, in *Climate Change 2007: The Physical Science Basis*, S. Solomon *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 2007).
- M. E. Conkright *et al.*, *World Ocean Atlas 2001*, vol. 4, *Nutrients*, S. Levitus, Ed., NOAA Atlas NESDros. Inf. Serv. 52 (U.S. Government Printing Office, Washington, DC, 2002).
- R. Schlitzer, *J. Phys. Oceanogr.* **37**, 259 (2007).
- We acknowledge the leadership of the Surface Ocean–Lower Atmosphere Study (SOLAS) project (www.solas-int.org) and the International Nitrogen Initiative (www.initrogen.org) of SCOPE and the International Geosphere-Biosphere Programme (IGBP; www.igbp.kva.se) for conceiving this synthesis. We thank the Scientific Committee on Oceanic Research, IGBP, the U.S. National Oceanic and Atmospheric Administration, and the European Science Foundation for partial support of the work. We thank J. Hare of the SOLAS International Project Office for help in organizing the workshop and E. Breviere of that office for help in making the Nitrogen Workshop in Norwich, UK, a success. We acknowledge two anonymous reviewers for constructive feedback.

Supporting Online Material

www.sciencemag.org/cgi/content/full/320/5878/893/DC1
SOM Text
Figs. S1 and S2
Table S1
References
10.1126/science.1150369