

Emerging patterns of marine nitrogen fixation

Jill A. Sohm, Eric A. Webb and Douglas G. Capone

Abstract | Biological N₂ fixation is an important part of the marine nitrogen cycle as it provides a source of new nitrogen that can support biological carbon export and sequestration. Research in the past decade has focused on determining the patterns of distribution and abundance of diazotrophs, defining the environmental features leading to these patterns and characterizing the factors that constrain marine N₂ fixation overall. In this Review, we describe how variations in the deposition of iron from dust to different ocean basins affects the limiting nutrient for N₂ fixation and the distribution of different diazotrophic species. However, many questions remain about marine N₂ fixation, including the role of temperature, fixed nitrogen species, CO₂ and physical forcing in controlling N₂ fixation, as well as the potential for heterotrophic N₂ fixation.

Upwelling

Wind-mediated movement of deep water to the surface.

Physical forcing

The effect of physical conditions and processes in the ocean on biological properties.

Diazotrophs

Organisms that can fix N₂ by converting it to ammonia.


Microbial N₂ fixation in the ocean is globally important because inputs of new N to surface waters (as opposed to N that is regenerated from existing organic material in surface waters) controls the export of organic matter to the deep ocean in many regions^{1,2}, a process commonly referred to as the ‘biological pump’ (REF. 3) (FIG. 1). Through the biological conversion of CO₂ to organic matter and its subsequent sinking, the deep ocean becomes a reservoir where CO₂ can be sequestered from the atmosphere for hundreds to thousands of years. Although it was initially acknowledged that N₂ fixation was a potential source of new N, the relative importance of newly fixed N was considered to be minor compared to the upwelling or diffusional flux of dissolved inorganic N (DIN) from deep waters². About a decade ago this view began to change, as it became apparent that marine N₂ fixation was a globally significant process that could equal the input of DIN to surface waters in areas of the tropical and subtropical ocean^{4,5}. Moreover, because upwelling and diffusion of DIN into surface waters brings with it an approximately stoichiometric amount of CO₂ (relative to the Redfield ratio (BOX 1)), N₂ fixation and atmospheric deposition of N are the only sources of new N that can lead to a net sequestration of atmospheric CO₂ in the deep ocean².

There are many factors that could limit and control marine N₂ fixation. In this Review, we specifically focus on the role of the nutrients P and Fe, as well as the potential effects of temperature, CO₂, fixed N and physical forcing on the distribution of N₂ fixation and diazotrophs in the ocean water column (although N₂ fixation also occurs in

various marine sedimentary environments; see BOX 2). We also highlight the potential of heterotrophic N₂ fixation in surface waters as an unrealized source of new N.

Nitrogen fixation — the basics

N₂ fixation is the process by which N₂ gas is reduced into two molecules of ammonia. The reaction consumes cellular energy, and the overall reaction is:



This reaction is catalysed by the nitrogenase enzyme complex, which is extremely oxygen sensitive. The most common form in the oceans is made up of two distinct proteins: dinitrogenase reductase (the Fe protein) and dinitrogenase (the Fe molybdenum, or FeMo protein)⁶. It is encoded by three genes (*nifH*, *nifD* and *nifK*); however, the *nif* operon can in some species contain up to 20 genes that are also involved in the synthesis and regulation of nitrogenase and its cofactors⁶. The nitrogenase proteins are highly similar among diazotrophs, and the well-conserved *nifH* gene is commonly used for phylogenetic and ecological studies⁶. Several ‘alternative’ nitrogenases are also known that contain a third subunit in the FeMo protein that is encoded by *nifG*⁶. The capacity for N₂ fixation occurs in bacteria and archaea of diverse physiologies (including anaerobes, facultative aerobes, aerobes and phototrophs), and although nitrogenase is thought to be an ancient enzyme, it is not uniformly distributed and is present in perhaps a few hundred cultivated species⁶.

Department of Biological Sciences and Wrigley Institute for Environmental Studies, University of Southern California, Los Angeles, California 90089, USA. Correspondence to J.A.S. e-mail: sohm@usc.edu doi:10.1038/nrmicro2594




Figure 1 | Overview of the importance of nitrogen fixation in the ocean. The export of organic material from surface into deeper waters is dependent on the amount of CO₂ fixation (primary production) that is based on new nitrogen inputs (N₂ fixation or nitrate diffusion from deep waters). Alternatively, some primary productivity is based on the regeneration of organic material to ammonia in surface waters. PON, particulate organic nitrogen.

The marine diazotrophs

In the marine water column, substantial N₂ fixation is carried out by several different types of diazotrophs, including non-heterocystous filamentous cyanobacteria, unicellular cyanobacteria and heterocystous filamentous cyanobacteria (both free-living and symbiotic forms), all of which are phototrophs but not necessarily autotrophs. Heterotrophic bacteria containing the *nifH* gene have also been found in the open ocean⁷, but their importance to global rates of N₂ fixation remains to be determined. The details of the physiology of marine diazotrophs have been presented in another recent review⁸, so they will be only briefly described here. In oligotrophic tropical and subtropical waters the non-heterocystous, filamentous diazotrophs are most commonly represented by the colony-forming cyanobacteria *Trichodesmium* spp.⁵. This genus has a unique physiology that employs spatial and temporal segregation and increased oxygen consumption to allow it to simultaneously fix N₂ and CO₂ (and thus evolve O₂) without a heterocyst⁹. *Trichodesmium* spp. grow and fix N optimally at temperatures between ~24 and 30 °C¹⁰ and are well known for forming large surface blooms when wind speeds are low and water temperature is above 25 °C⁵. The collapse of these blooms could contribute to organic matter export.

The occurrence of N₂ fixation and the diversity of diazotroph species in the <10 μm plankton fraction was only recognized about a decade ago¹¹. These unicellular diazotrophs are divided into three groups: *Crocospaera watsonii*, a cultured cyanobacterium that has many representative strains available; UCYN-A, an uncultured cyanobacterium; and UCYN-C, a distinct, uncultured clade that is related to *Cyanothece* sp. ATCC

51142 (REF. 8). They are commonly classified by group-specific quantitative PCR (qPCR) of the *nifH* gene in DNA samples, in which *nifH* copy number is assumed to equal the cell number of the specific diazotroph. These three groups of small diazotrophs have been found in diverse tropical and subtropical surface waters^{12,13}, and their N₂ fixation rates can equal or exceed that of *Trichodesmium* spp. when they are present at high densities^{14,15}.

C. watsonii carries out photosynthetic CO₂ fixation during the day and N₂ fixation at night⁸. Despite the fact that strains of this species are extremely similar phylogenetically^{16,17}, the isolates fall into two distinct phenotypic groups: large (>4 μm) and small (<4 μm) cells. The large isolates display greater rates of N₂ fixation than the small isolates¹⁶ and also produce large amounts of extracellular polysaccharides (J.A.S., E.A.W., B. R. Edwards and B. G. Wilson, unpublished). The temperature range for growth of *C. watsonii* is ~22–36 °C¹⁶. In the field, DNA copy numbers of *nifH* from *C. watsonii* increase linearly with temperature, reaching a maximum of nearly 10⁶ *nifH* copies per litre at temperatures of 30 °C¹⁸, but are generally closer to 10³–10⁴ per litre^{12,13}. Because UCYN-A remains uncultured, its physiology has been inferred from a recent analysis of an environmental metagenome. The UCYN-A genome contains photosystem I (PSI) genes but lacks genes for photosynthetic accessory pigments, the oxygen evolving photosystem II (PSII), the tricarboxylic acid cycle and metabolic pathways for some amino acids and purines, indicating that UCYN-A is photoheterotrophic and possibly lives in association with other phototrophs^{19,20}. In the ocean, *nifH* DNA has been amplified from UCYN-A that are found at temperatures ranging from 15–30 °C^{12,13,21}, with the greatest copy number being found at ~24 °C¹⁸, indicating that this group has a wider temperature range and lower minimum temperature than the other marine diazotrophs. UCYN-A bacteria are often the more abundant of the unicellular diazotrophs in oligotrophic waters, ranging from 10³–10⁵ *nifH* DNA copies per litre on average, and sometimes reaching abundances of 10⁶ copies per litre^{12,18}.

The heterocystous diazotrophs are a filamentous group that includes both free-living and symbiotic cell types, which are typically restricted to more specialized marine environments. The cooler, brackish waters of the Baltic Sea host *Nodularia* spp., *Aphanizomenon* spp. and sometimes *Anabaena* spp. The temperature optima of *Aphanizomenon* spp. and *Nodularia* spp. isolates are 16–22 °C and 25–28 °C, respectively²²; both are present at around 10⁴ heterocysts per litre in their respective blooms²³. In warmer waters of the tropics and subtropics, heterocystous diazotrophs (such as *Richelia intracellularis*) are only encountered as symbionts of diatoms like *Hemiaulus* spp., and *Rhizosolenia* spp. Filaments of *Calothrix* spp. can also be found in association with the diatom *Chaetoceros* spp.²⁴. These diatom diazotroph associations (DDAs) are widely distributed through the warm oligotrophic ocean^{25–27}, averaging 10³ heterocysts per litre in a bloom, but sometimes reaching 10⁶ heterocysts per litre²⁶. Because

Heterocyst

A specialized cell with a thick cell wall that lacks photosystem II (PSII) and is the site of N₂ fixation in some filamentous cyanobacteria.

Blooms

Areas of large growth or accumulation of a species.

Brackish

Describing water that has a salinity between fresh and marine water.

Box 1 | The Redfield ratio

More than half a century ago, Alfred C. Redfield¹²¹ recognized that the dissolved nutrient ratios in the deep sea corresponded remarkably to the ratios of particulate organic matter in surface waters. He attributed this correlation to the fact that organic matter (phytoplankton biomass) is produced, on average, in a ratio that is determined by biochemical needs, and that this material is subsequently remineralized in the deep ocean, releasing inorganic nutrients in the same proportions. This ratio is now referred to as the Redfield ratio, which has been determined to be 106C/16N/1P. In the marine environment, the ratio of N/P can vary substantially in dissolved forms in surface waters, and also in particulate matter, and the Redfield ratio is often cited as the line between N and P limitation, with higher ratios indicating P limitation and lower ratios indicating N limitation. It is important to acknowledge that many organisms have N/P ratios that differ from 16/1; however, the uniformity of the average deep water dissolved inorganic N/dissolved inorganic P (DIN/DIP) ratio around the globe strongly supports an average phytoplankton demand of 16/1.

of their heavy, silicon-containing cell walls, DDAs may have an important role in C export²⁸, and could therefore be an important part of the biological pump. Other planktonic marine diazotrophs that have been identified by *nifH* gene diversity assessments are members of the gammaproteobacteria and alphaproteobacteria, and members of the anaerobic cluster III diazotrophs^{7,11,13}. The importance of these groups to open ocean marine N₂ fixation is currently unknown and represents an exciting area of future research.

Iron and phosphorus limitation

Marine phytoplankton are generally thought to be N limited through much of the tropical and subtropical oceans²⁹, whereas diazotrophs by definition are not. This suggests that other nutrients control diazotrophs in the ocean, with the most likely candidates being Fe and P. The importance of each of these nutrients is probably determined by the prevailing inputs and cycling of nutrients in the different ocean basins (see BOX 3 for background information).

Iron limitation of diazotrophs in culture. Fe limitation may be important for cyanobacterial N₂ fixation as Fe is a cofactor in PSI and nitrogenase (as well as in other redox enzymes), such that the Fe requirement for diazotrophy is ~5 times higher than that for ammonia assimilation^{30,31}. The concentration of dissolved Fe (dFe) is also low in many of the environments where N₂ fixation occurs^{32–35}. Early work established the importance of Fe to N₂ and CO₂ fixation and growth in natural populations and cultures^{36,37}, and recent research has probed the mechanisms behind this importance. Low free Fe concentrations decrease N₂ fixation rates in multiple strains of *Trichodesmium* spp.^{31,38}, causing the expression of Fe stress genes that are predicted to compensate for Fe deprivation, and downregulation of Fe-containing proteins that are involved in photosynthesis and N₂ fixation^{38–40}. The expression of these genes is not necessarily specific to diazotrophs; however, group-specific expression of these genes can be used to detect Fe stress in the field. The observed decrease in N₂ fixation may be due to a decrease in nitrogenase protein levels under conditions of Fe limitation, while the cell maintains the

abundance of the Fe-requiring PSI³⁹. N₂ fixation and growth in *C. watsonii* are also reduced under conditions of Fe depletion⁴¹. However, to reduce its overall Fe needs, *C. watsonii* synthesizes Fe-containing proteins for N₂ fixation and CO₂ fixation only during the night or day, respectively, repurposing the Fe throughout the diel cycle and thus reducing the total cellular Fe requirement by 40%⁴². Fe deficiency can also reduce the growth and yield in *Nodularia spumigina*; however, *N. spumigina* may be able to overcome some of this effect by accessing ligand-bound Fe⁴³.

Phosphorus limitation of diazotrophs in culture. P is only supplied to the ocean by geological processes (that is, weathering) and is a major component of cells, having roles in information storage (DNA and RNA), protein synthesis (ribosomal RNA), energy generation (ATP) and as a component of cell membranes (phospholipids). Despite the fact that *Trichodesmium* spp. and other phytoplankton can substitute sulpholipids for phospholipids during P deficiency⁴⁴, they still are not released from a requirement for P for growth.

Trichodesmium spp. culture work has identified a number of mechanisms to deal with P limitation. Maximal dissolved inorganic P (DIP) uptake rates⁴⁵, alkaline phosphatase activity (APA)⁴⁶ and particulate N/P ratios^{47,48}, as well as the transcription of genes for DIP transporters and alkaline phosphatases⁴⁹, all increase under conditions of DIP-induced limitation of growth rate. This allows *Trichodesmium* spp. to deal with P limitation through a greater uptake of DIP and dissolved organic P (DOP), and through decreasing the requirement for cellular P. These physiological and gene expression measures can be used as markers for P stress in the field. Additionally, *Trichodesmium erythraeum* str. IMS101 possesses the full *phn* operon, which gives it the genetic capability to use phosphonate (C–P) bonds⁵⁰. Other isolates of *Trichodesmium* spp. contain the *phnJ* and *phnD* genes, suggesting that the *phn* genes are common in the genus. In culture and field populations, expression of *phnJ*, which is thought to encode part of a phosphonate bond lyase complex, was detected when cells were DIP deficient⁵⁰. Direct uptake of P from phosphonates has yet to be shown, but *Trichodesmium* spp. cultures seem to be able to grow with phosphonate bond-containing compounds as the sole P source⁵¹. Despite the fact that *Trichodesmium* spp. have many physiological adaptations for dealing with P limitation, N₂ fixation can still be reduced in cultures that are depleted of DIP⁴⁷. Similarly, DIP deficiency in *C. watsonii* leads to the expression of a high-affinity phosphate transporter, but only the yield and not the growth rate of cultures is reduced⁵². *C. watsonii* isolates can also grow on a number of DOP compounds containing phosphomonoesters⁵² (but not phosphonates). P deficiency can also affect the growth of Baltic Sea isolates of *Aphanizomenon* spp. and *Nodularia* spp. Both species decrease their requirement for P and growth rates under DIP deficiency; however, *Aphanizomenon* spp. seem to be adapted to take advantage of DIP pulses, whereas *Nodularia* spp. have adaptations for growing well at low DIP concentrations⁵³.

Dissolved Fe

Fe that can pass through a 0.4 µm filter.

Free Fe

A pool of Fe that is composed of Fe bound to inorganic ligands and a small amount of free ion (Fe³⁺) and is presumed to be bioavailable (often denoted Fe⁰).

Ligand

In this article, an organic molecule that binds Fe (potentially a siderophore).

Sulpholipid

A membrane lipid containing a sulphur (as opposed to P)-based head.

Alkaline phosphatase

A hydrolytic enzyme that cleaves phosphomonoesters from P-containing dissolved organic matter.

Box 2 | Emerging patterns in benthic marine nitrogen fixation

In addition to occurring in the pelagic zone, N_2 fixation occurs in benthic habitats such as microbial mats and plant rhizospheres. Research in these areas was brisk in the 1980s and before^{122,123}, but few new discoveries were made in subsequent years. However, in the past 5 years interest in benthic marine N_2 fixation has been renewed and researchers are again turning to the sediments. Recent research has shown that N_2 fixation might be an important process in estuaries, with N_2 flux shifting into the sediment in Narragansett Bay in 2006, indicating net N_2 fixation¹²⁴. N_2 fixation has also been detected in areas that were not typically studied in the past, such as bare sediments (sediments free of visible microbial mats), depths >5 cm¹²⁵ and the deep sea^{126,127}. In shallow marine sediments, N_2 fixation can be attributed almost entirely to sulphate-reducing bacteria¹²⁵, and this activity correlates with macrofauna behaviour. In the deep sea, a microbial consortium has been identified that contains an archaeon and bacterium that together carry out the anaerobic oxidation of methane. The archaeon also appears to have the ability to fix N_2 and share it with its associated bacterium¹²⁶. Molecular methods are also driving research into diel patterns of N_2 fixation by specific cyanobacteria and heterotrophic bacteria¹²⁸. Still, there are many more questions left to be answered; for example, why do microorganisms fix N_2 in sediments when reduced nitrogen species are in abundance in porewaters?

Iron and phosphorus limitation in the Atlantic Ocean.

Early work by Rueter³⁷ showed that the introduction of Fe increases CO_2 and N_2 fixation rates and cellular chlorophyll *a* content in natural samples of *Trichodesmium* spp. that were collected near Barbados. Aside from this study and a more recent study showing P and Fe co-limitation of bulk N_2 fixation at two stations in the eastern tropical North Atlantic Ocean⁵⁴, Fe does not seem to be an important factor in limiting the rate of N_2 fixation (in the sense of Blackman limitation) in the North Atlantic Ocean. However, it appears that the amount of Fe delivered in dust is important in determining the diazotrophic biomass (in the sense of Liebig limitation) in this basin⁵⁵, and that high rates of N_2 fixation ($250 \mu M m^{-2} d^{-1}$ from *Trichodesmium* spp. alone (BOX 3)) in turn drive the North Atlantic Ocean towards P limitation, with DIP concentrations as low as $1 nM$ ^{56,57} (FIG. 2). P stress in *Trichodesmium* spp. is suggested by the maximal DIP uptake rates, levels of APA and the particulate N/P ratios from the North Atlantic Ocean and the South Atlantic Ocean offshore of Brazil^{47,58,59}. Furthermore, N_2 fixation by *Trichodesmium* spp. is proportional to P requirements in colonies in the western North Atlantic Ocean⁶⁰. These data are consistent with recent research showing a higher degree of P stress in *Trichodesmium* spp. in the western versus the eastern North Atlantic Ocean⁶¹. Across the South Atlantic Ocean gyre, surface concentrations of DIP are one to two orders of magnitude greater than in the North Atlantic Ocean, whereas dFe is ~3 times lower⁵⁵, indicating that Fe may be limiting to N_2 fixation in the South Atlantic Ocean. Indeed, dFe may be so low as to preclude N_2 fixation, as rates are $\leq 20 \mu M m^{-2} d^{-1}$ (REF. 55) (BOX 3).

Iron and phosphorus limitation in the Pacific Ocean.

Based on observational data of dust inputs and nutrient concentrations, many researchers have suggested that Fe limits N_2 fixation in the Pacific^{35,57,62,63}. Subsequent field work has shown that addition of extra Fe can substantially increase N_2 fixation rates in the North Pacific at some times of the year⁶⁴, and that *Trichodesmium* spp. express

an Fe stress gene over much of the South Pacific Ocean (P. D. Chappell, J. W. Moffett, A. M. Hynes and E.A.W., unpublished). Additionally, dust inputs seem to drive N_2 fixation in waters near Japan⁶⁵. A decrease observed over a 10 year period in DIP stocks at station ALOHA (an autonomous ocean observatory in the North Pacific Ocean near Hawaii) is thought to be linked to the rise of N_2 fixing cyanobacteria and suggests a shift in this system to P limitation⁶⁶. More recent data suggest that although P is becoming scarcer in the North Pacific Ocean, diazotrophs living there are not P limited^{58,67,68} (FIG. 2). In the South Pacific Ocean gyre, dFe concentrations are some of the lowest that have been measured in the ocean: 0.1–0.2 nM in open ocean areas³³. The extremely low DIN (<0.02 μM)³³ and DIP concentrations that are around an order of magnitude greater than in the North Atlantic Ocean⁶⁹ (and similar to the South Atlantic Ocean) would seem to favour N_2 fixation. However, N_2 fixation rates observed in the region are $\sim 50 \mu M m^{-2} d^{-1}$ (REF. 70) (BOX 3), leading to the hypothesis that the low dFe concentrations limit N_2 fixation in the South Pacific Ocean.

Iron and phosphorus limitation in the Baltic Sea.

The Baltic Sea is a brackish inland sea that is known to host large blooms of *Nodularia* spp., *Aphanizomenon* spp. and *Anabaena* spp. during the summertime. In coastal zones, Fe can stimulate diazotroph biomass⁷¹, but in the open sea of the Baltic, research has converged on the importance of P to diazotrophic activity^{72–74}. A number of studies have linked the strength of diazotroph blooms to the amount of excess DIP left over after the spring bloom^{72,75}, although another study suggested that an unidentified P source must also be important, probably DOP⁷⁶. Nevertheless, the cyanobacterial blooms perform luxury uptake of DIP in spring, exhibiting particulate N/P ratios of ~5. As the bloom grows, N/P ratios increase to ~80 before bloom collapse⁷⁷. These blooms induce a transient P limitation in the Baltic Sea microbial community during the summer^{74,78} and also the limitation of N_2 fixation itself^{73,78}. Interestingly, the *Aphanizomenon* spp. bloom generally initiates a few weeks before the numbers of *Nodularia* spp. increase, and the intensity of the *Aphanizomenon* spp. bloom can be linked to the amount of excess DIP⁷⁵. *Nodularia* spp. seem to respond more to temperature and may be able to use regenerated P⁷⁵, a strategy that explains the delay in blooming of this species compared to *Aphanizomenon* spp.

Iron and phosphorus limitation in other areas.

Relatively limited information is available for other bodies of water. The Mediterranean Sea is thought to be severely P limited⁷⁹ owing to high DIN/DIP ratios⁸⁰ that also point to the Mediterranean Sea as a site of N_2 fixation. Conversely, high N/P ratios of nutrient inputs to the Mediterranean Sea may account for the high DIN/DIP ratios. Despite this suggestion, there have been few N_2 fixation studies in the region. Recent work has identified putative diazotrophs from the cyanobacteria, proteobacteria, cluster III and methanogenic archaea groups⁸¹. Microscopy methods have been used to show that diazotrophic cyanobacteria are generally found free-living in

Blackman limitation
The limitation of the growth rate of an organism.

Liebig limitation
The limitation of the yield of an organism (as in a crop yield).

Gyre
A giant circular surface current that is present in the ocean.

Luxury uptake
The uptake of nutrients that are in excess of demand, and that can be made into storage products.

Box 3 | Setting the stage: the biogeochemistry of the ocean

N_2 fixation is generally thought to occur in warm, oligotrophic waters stretching roughly from 30°N to 30°S (REF. 5). Fe in the open ocean derives from the deposition of dust from adjacent desert areas. In general, the oceans in the northern hemisphere receive greater dust inputs than the southern oceans owing to the greater land mass in the northern hemisphere⁶³. This is particularly true in the North Atlantic Ocean, where the Sahel and Sahara provide a large dust source causing dissolved Fe (dFe) concentrations of around 1 nM, compared to ~0.3 nM in the South Atlantic Ocean^{22,55} (see the table). dFe levels are consistent across the entire Pacific at 0.1–0.2 nM^{33,62}. However, dFe is elevated near islands such as the Hawaiian archipelago⁶². In the Arabian, Baltic and Mediterranean Seas dFe concentrations are relatively high, ≥ 1 nM^{129–131}, with concentrations in the Baltic Sea probably being affected by land runoff^{71,131}.

Dissolved inorganic P (DIP) concentrations in much of the open ocean range from 100–200 nM^{55,69,132} except in the North Atlantic, where they can be below 1 nM in the western basin^{56,57}. Dissolved inorganic nitrogen (DIN) concentrations, conversely, are universally low in oligotrophic surface waters^{33,55,132}, causing nutrient ratios in the North and South Pacific Oceans, and the South Atlantic Ocean to be much lower than the Redfield ratio, and much higher than 16/1 in the North Atlantic. DIP concentrations in the Arabian and Baltic Seas are seasonally variable and can be >1 μ M during mixing periods^{72,86}, but undetectable in the warm stratified period, with DIN/DIP ratios of less than 16/1 (REFS 78,86). DIP concentrations are also extremely low in the Mediterranean Sea owing to the anti-estuarine circulation of water, and DIN/DIP ratios are consequently high at ~30 (REF. 80).

Because of differences in sampling effort, N_2 fixation rates in some areas are better understood than others. In the North Atlantic Ocean, N_2 fixation by *Trichodesmium* spp. alone can be 250 μ mol N m^{-2} d^{-1} in the season in which N_2 fixation occurs⁴, whereas bulk water rates are only about 20 μ mol N m^{-2} d^{-1} in the South Atlantic Ocean⁵⁵. In the North Pacific Ocean, rates are about 100 μ mol N m^{-2} d^{-1} (REFS 15, 133) but can be much higher, and rates in the South Pacific Ocean are 50 μ mol N m^{-2} d^{-1} (REF. 70). Biological estimates of N_2 fixation in the Arabian Sea range from 16–122 μ mol N m^{-2} d^{-1} (REF. 85). N_2 fixation in the Baltic Sea is highly seasonal, ranging from as low as 20 μ mol N m^{-2} d^{-1} in the winter up to 2,500 μ mol N m^{-2} d^{-1} during summertime cyanobacterial blooms¹³⁴, whereas rates seem extremely low in the Mediterranean Sea at ~3 μ mol N m^{-2} d^{-1} (REF. 83).

Location	Dust deposition	dFe (nM)	DIP (nM)	DIN/DIP	N_2 fixation (μ mol N m^{-2} d^{-1})
North Atlantic Ocean	very high (REF. 63)	0.2–1.2 (REFS 32,55)	0.2–5 (REFS 56,57)	~30 (REF. 57)	250 (annual average) (REF. 4)
South Atlantic Ocean	very low (REF. 63)	0.1–0.4 (REF. 55)	~200 (REF. 55)	~0.1 (REF. 55)	~20 (spring) (REF. 55)
North Pacific Ocean	low (REF. 63)	0.1–0.3 (REF. 62)	10–100 (REF. 132)	~1 (REF. 57)	30–120 (annual range) (REF. 133)
South Pacific Ocean	very low (REF. 63)	0.1–0.2 (REF. 33)	110–240 (REF. 69)	~0.1 (REF. 33)	30–70 (spring) (REF. 70)
Arabian Sea	high (REF. 63)	0.5–1 (REF. 129)	BD–1,000 (REF. 86)	>13 (REF. 86)	16–122 (annual estimate) (REF. 85)
Mediterranean Sea	high (REF. 63)	0.2–1.2 (REF. 130)	BD–70 (REF. 135)	~28 (REF. 80)	~3 (spring and summer) (REF. 83)
Baltic Sea	low (REF. 63)	3–7 (REF. 131)	BD–800 (REF. 72)	5–40	~2500 (summer peak) (REF. 134)

BD, below detectable limits.

low abundance and ranging in size from 0.2–3 μ m⁸², and DDAs are also found in low abundance²⁵. Measured N_2 fixation rates are also very low (~3 μ M m^{-2} d^{-1})⁸³ (BOX 3), and it was recently concluded that extreme P limitation precludes substantial N_2 fixation in the eastern basin of the Mediterranean Sea⁸⁴.

The Arabian Sea has long been known to host large blooms of *Trichodesmium* spp.⁸⁵; however, the constraints on N_2 fixation rates in this region are not well understood. The presence of large amounts of excess DIP in surface waters⁸⁶ and the apparent link between *Trichodesmium* spp. abundances and dust deposition⁸⁷ suggest that Fe may be the limiting nutrient. This area is also strongly affected by physical factors, and blooms may be triggered by the calm, inter-monsoon periods⁸⁷. In the nearby Red Sea, *Trichodesmium* spp. and gammaproteobacteria have been detected, and N_2 fixation rates are low and possibly P limited⁸⁸.

Diazotrophic distribution in the open ocean

The distribution of physiologically and morphologically different diazotrophic species has ramifications for the potential for N_2 fixation in a given area and the fate of recently fixed N in surface waters. It is apparent from qPCR quantification and microscopy techniques that *Trichodesmium* spp. are the most abundant and perhaps the dominant diazotrophs in parts of the North Atlantic Ocean^{12–14,89}, whereas the Pacific can be dominated — at least numerically — by the unicellular diazotrophs, particularly UCYN-A^{12,18} (FIG. 2). Recent work indicates that dFe concentrations could be controlling the distribution of diazotrophic species. *Trichodesmium* spp. are abundant in the North Atlantic Ocean, where dFe concentrations are relatively high, but not in the South Atlantic Ocean, where dFe concentrations are extremely low⁵⁵. Additionally, *Trichodesmium* spp. may have a selective advantage in low P areas (that is, the North Atlantic




Figure 2 | Summary of the distribution of nitrogen fixation rates, diazotrophic species and nutrient limitation in the ocean. Relative N₂ fixation and information on the prevailing limiting nutrient are shown in each area by the fill colour and outline, respectively, and the source and relative strength of dust sources are shown by the size of the grey arrows. The diazotrophic species that dominate in each basin are indicated, with other types noted if they are found at high abundance in specific environments. Much of the abundance data available come from late spring–early autumn in each hemisphere, when higher overall abundances are expected. Robust seasonal data sets of diazotroph biomass are only available for limited locations, so it is less clear how the global patterns may change temporally. DDA, diatom diazotroph association; EQ, equator.

Ocean), as they have the genetic capability to use phosphonates. Within each basin, there is also evidence for smaller-scale variation in the populations. In one study in the tropical North Atlantic Ocean, *Trichodesmium* spp. were shown to dominate across much of the basin, except near the Cape Verde Islands where UCYN-A was most abundant¹⁴. In the Pacific, *Trichodesmium* spp. can be seen in relatively high abundances near the Hawaiian Islands⁹⁰, where dFe concentrations are elevated^{34,62}. In the central gyre, however, dFe concentrations are very low (~0.1 nM)⁶², *Trichodesmium* spp. are not common, and N₂ fixation seems to be carried out by small unicellular cyanobacteria^{12,90}. The apparent dominance of the North Pacific Ocean by UCYN-A¹² supports the hypothesis that the extremely low Fe concentrations inhibit *Trichodesmium* spp. growth in the North Pacific Ocean, but are high enough for picodiazotrophs and nanodiazotrophs. A similar distribution of N₂ fixation is seen in the equatorial Pacific, with fixation being carried out by *Trichodesmium* spp. near Papua New Guinea but by small diazotrophs in open waters⁹¹. In the understudied central southern gyres of the Atlantic and Pacific Oceans, there is evidence of the occurrence of unicellular cyanobacteria¹⁸, but more work needs to be done to confirm the importance of unicellular diazotrophs in this region.

Heterocystous symbionts have a much patchier distribution throughout the world's oceans, probably because the diatom hosts require silicon to build their cell walls. Hence, research to date has identified the largest densities of DDAs in the Amazon River plume^{14,26,28,92}, which delivers high concentrations of nutrients — including silicate — to a large area of the tropical North Atlantic Ocean²⁸. DDAs have also been found in the Gulf of Guinea near the Congo River plume⁹³, the Mediterranean Sea²⁵ and across the Pacific Ocean²⁷. In these regions, abundance is low compared to the Amazon River plume, and it is not clear what promotes the growth of these organisms in the open ocean. Although their distribution and abundance appear patchy, evidence suggests that DDAs are highly efficient at exporting C (REF. 28), and thus these species could be very important where they occur.

Given the differential dominance of different groups of diazotrophs in the oceans, it is likely that the fate of recently fixed N will also differ geographically. The modes of transfer of newly fixed N into the food web or of export from surface waters are not well known, but probably differ based on morphology, chemical composition and size differences between diazotrophic species⁶. *Trichodesmium* spp. have only one well

characterized grazer, *Macrosetella gracilis*, which lives its life attached to *Trichodesmium* spp. colonies and is probably only responsible for a small amount of mortality⁹⁴. However, *Trichodesmium* spp. also appear to release newly fixed N⁹⁵. To date, nothing is known about the grazing, recycling or export of *C. watsonii* and UCYN-A. Grazing, sinking and nutrient release in diazotrophs control the fate of newly fixed N and thus its impact on the global C cycle. These factors are important to consider in the future.

Future study of marine nitrogen fixation

Nitrogen fixation in lower temperature waters. Although marine N₂ fixation is generally associated with warmer tropical and subtropical surface waters, the process of N₂ fixation is not intrinsically inhibited by temperature and can occur at temperatures near 0 °C⁹⁶. However, individual species have optimal temperatures, thus temperature may affect species distribution; for example, it has been proposed that warm waters allow for the high respiration rates that preclude the need for a heterocyst and thus select against free-living heterocystous forms⁹⁷. Colder surface waters at higher latitudes are often associated with upwelling of nutrients (including fixed N), presumably selecting against N₂ fixation. Despite this, N₂ fixation by small diazotrophs has been reported in two Leuwin Current eddies at temperatures of 18 °C and 15 °C⁹⁸. It has also been found at 18–20 °C in the English Channel (which was linked through an mRNA *nifH* library to the presence of UCYN-A, proteobacteria and the anaerobic cluster III group⁹⁹) and at 19 °C off the coast of California (which was attributed to UCYN-A²¹). Thus, N₂ fixation by small diazotrophs can occur at temperatures that are lower than previously thought, but deeper investigation of the physiological basis of these occurrences is warranted.

Control of nitrogen fixation by carbon dioxide. In light of increasing atmospheric CO₂ concentrations and acidification of surface waters, it is important to understand how these related factors will affect N₂ fixation in the ocean in the future. Experiments by multiple investigators with *T. erythraeum* str. IMS101 show higher growth and N₂ fixation rates under increased CO₂ concentrations^{100–102}, suggesting that N₂ fixation by open ocean diazotrophs could provide negative feedback on the increase in global CO₂ concentrations through an increase in new N to drive the biological pump. However, the positive effect of increased CO₂ at current ocean temperatures is not as pronounced when temperatures are increased¹⁰³. *C. watsonii* str. WH8501 shows a similar response to *T. erythraeum* str. IMS101 when faced with increased CO₂⁴¹; however, CO₂ had no effect during Fe depletion, indicating that low Fe could temper the effect of CO₂ on *C. watsonii* in future climate scenarios. Higher CO₂ concentrations may allow greater efficiency in the C concentrating mechanism, freeing up energy to be shunted towards N₂ fixation^{100–102,104}. In *N. spumegina*, increased CO₂ may have the opposite effect, substantially decreasing growth and N₂ fixation rates¹⁰⁵. The authors speculate that more acidic conditions impede

the transfer of compounds between heterocystous and vegetative cells¹⁰⁵. However, more research on this topic is needed before general conclusions can be drawn.

Research on the interaction of factors that are likely to change in concert and the long-term effects of exposure to higher CO₂ are the next step to truly understanding how N₂ fixation will change in the future in a human-influenced ocean.

Fixed nitrogen as an inhibitor of nitrogen fixation.

Fixed N has long been recognized as an inhibitor of N₂ fixation, as it is an energetically costly process to carry out in comparison to ammonia assimilation. N₂ fixation in most diazotrophs examined is inhibited by mM concentrations of ammonia¹⁰⁶, whereas mM concentrations of nitrate shut down N₂ fixation in some strains¹⁰⁷ but have little to no effect in others¹⁰⁸. The increased energy requirement for the reduction of nitrate after uptake compared to 'direct' assimilation of ammonia might explain these differences. A recent modelling exercise predicted that some of the largest areas of N₂ fixation in the ocean occur in or near waters that contain measurable amounts of nitrate¹⁰⁹, thus some recent studies have begun to question the strength of feedback inhibition of fixed N species on marine N₂ fixation. Addition of ammonia to natural samples of *Trichodesmium* spp. can shut down nitrogenase activity¹¹⁰; however, culture studies have shown mixed results, with some indicating that short-term additions of fixed N sources do not reduce activity^{111,112}, whereas others show a reduction in activity under these conditions¹¹³. Finally, N₂ fixation has been shown in the nitrate-rich upwelling areas off Chile⁷⁰, demonstrating that N₂ fixation may not be as strongly regulated by fixed N as previously presumed.

Physical forcing of N₂ fixation. It has long been known that biological processes in the ocean can be driven by physical processes, but only recently have researchers specifically begun to look at physical forcing of N₂ fixation. Davis and McGillicuddy¹¹⁴ found that *Trichodesmium* spp. abundance in the North Atlantic Ocean correlated with anticyclonic eddies, suggesting that N₂ fixation is enhanced inside these mesoscale features. This phenomenon has since been studied in the North Pacific Ocean, where N₂ fixation in an anticyclonic eddy was found to be higher than the rates that are typically found at nearby Station ALOHA¹¹⁵. Over a 3-year time-course, spikes in N₂ fixation at Station ALOHA can also be linked to sea surface height anomalies that suggest the presence of anticyclonic eddies¹¹⁶. Finally, depth-integrated N₂ fixation was found to be 2.5 times greater in a warm core eddy versus a cold core eddy off the west coast of Australia⁹⁸. These studies provide intriguing insights, but more research is needed to fully understand the mechanisms behind physical forcing of N₂ fixation in the ocean.

Heterotrophic nitrogen fixation. Heterotrophic N₂ fixation is a potentially new paradigm in marine N₂ fixation, as heterotrophic organisms are secondary producers that use dissolved organic C for energy generation. This

Eddies

Small-scale circular currents in which the water inside is typically different from the water outside.

C concentrating mechanism

The mechanism that increases the concentration of CO₂ around ribulose-1,5-bisphosphate carboxylase oxygenase (RuBisCo), in order to increase the rate of photosynthesis.

Anticyclonic eddies

Eddies with warm interior water and a centre that is slightly higher than the surrounding sea surface. They rotate clockwise in the northern hemisphere and anticlockwise in the southern hemisphere.

is an important difference from the more commonly recognized primary producing diazotrophs discussed above (not including UCYN-A, which is photoheterotrophic), as they will not concomitantly fix CO₂, but will instead produce it. Throughout the world's oceans, non-cyanobacterial *nifH* genes have been identified including those of alphaproteobacteria, gammaproteobacteria and phylotypes from *nifH* of cluster III (see REF. 7); however, it is unclear whether this genetic potential translates into nitrogenase activity. Evidence of expression of these *nifH* genes has been found in cDNA libraries from the Mediterranean Sea⁸¹, the eastern North Atlantic Ocean and the English Channel⁹⁹, and qPCR has been used to quantify expression of *nifH* in gammaproteobacteria^{117,118}. However, direct uptake of N₂ by heterotrophic bacteria in the water column has yet to be shown, thus the potential contribution of these organisms to marine N₂ fixation is undetermined. Additionally, how do organic C availability and concentration control the process of N₂ fixation, and what mechanism(s) do diazotrophs use to protect nitrogenase from O₂ inactivation? These questions highlight only a few reasons why research on heterotrophic N₂ fixation is likely to gain momentum in the coming years. We speculate that the attachment of bacteria to particles could be a mechanism to deal with some of the challenges heterotrophic diazotrophs might face.

The emerging global picture

The accumulating data suggest a range of factors that may constrain N₂ fixation *in situ*. On the larger scale, temperature seems to set the limits of where different diazotrophs can exist, although it is important to note that temperature may be correlated to other factors that control geographical ranges. Large dust inputs into the tropical North Atlantic Ocean from the Sahel and Sahara appear to foster higher availability of Fe in the tropical and subtropical North Atlantic Ocean, thereby stimulating N₂ fixation, which causes drawdown of DIP and makes DIP availability a more important constraint on N₂ fixation than Fe. By contrast, excess DIP in surface waters of the tropical and subtropical Pacific Ocean (both North and South) and the South Atlantic Ocean coupled with low deposition rates of Fe from the atmosphere seems to drive diazotrophy to Fe limitation⁵⁵. Recent modelling efforts have integrated physiological parameters of diazotrophs and generated global maps of diazotroph limitation that

are roughly consistent with this picture¹¹⁹, suggesting that the underlying controls on diazotrophy are becoming better understood. However, it is important to note that high or low dust deposition does not necessarily predict the presence of N₂ fixation. For example, relatively high dust deposition to the Mediterranean Sea should support N₂ fixation there except DIP levels are too low, whereas N₂ fixation occurs at extremely high rates in the landlocked Baltic Sea despite the relatively low dust deposition (Fe and other nutrients are probably supplied by river and terrestrial runoff). The presence of fixed N probably also has an effect on global diazotroph distributions by selecting against N₂ fixation; however, the concentration at which inhibition occurs may be higher than previously thought and dependent on the type of diazotrophs present. Finally, at current atmospheric concentrations, CO₂ may also have a role in determining the total amount of N₂ fixation in the ocean and might increase N₂ fixation rates in the future as humans continue to discharge CO₂ into the atmosphere.

In parallel, recent molecular evidence indicates that the relative dominance of different diazotrophic groups also varies among ocean basins. Microdiazotrophs such as *Trichodesmium* spp. seem to have a more predominant role in the North Atlantic Ocean compared to nanodiazotrophs such as *C. watsonii* and other small coccoid cyanobacteria^{13,92}, whereas nanodiazotrophs are seemingly more dominant though much of the North Pacific Ocean^{12,91}. We posit that the cause of these emergent patterns is the differences in nutrient inputs and cycling in distinct parts of the ocean (FIG. 2). Understanding the distribution of these diazotrophic species is important because the fate of the N (and potentially C) fixed by these different and contrasting groups may follow distinct pathways into marine food webs, thus differentially influencing the biological pump.

With the prospect of upper ocean warming and acidification as a result of global climate change, as well as the increasing anthropogenic flux of reactive N to the sea¹²⁰, the magnitude, distribution and importance of marine N₂ fixation is likely to change, although research into how is just beginning. Despite decades of advances, the marine N cycle remains poorly understood, and with climate- and human-induced changes on the horizon, the continued study of oceanic N₂ fixation is important for our broader understanding of ocean biogeochemistry now and in the future.

- Dugdale, R. C. & Goering, J. J. Uptake of new and regenerated forms of nitrogen in primary productivity. *Limnol. Oceanogr.* **12**, 196–206 (1967).
- Eppley, R. W. & Peterson, B. J. Particulate organic-matter flux and planktonic new production in the deep ocean. *Nature* **282**, 677–680 (1979).
- Longhurst, A. R. & Harrison, W. G. The biological pump – profiles of plankton production and consumption in the upper ocean. *Prog. Oceanogr.* **22**, 47–123 (1989).
- Capone, D. G. *et al.* Nitrogen fixation by *Trichodesmium* spp.: an important source of new nitrogen to the tropical and subtropical North Atlantic Ocean. *Global Biogeochem. Cycles* **19**, GB2024 (2005). **An extensive review of the physiology, ecology and biogeochemistry of *Trichodesmium* spp. in the North Atlantic Ocean.**
- Capone, D. G., Zehr, J. P., Paerl, H. W., Bergman, B. & Carpenter, E. J. *Trichodesmium*, a globally significant marine cyanobacterium. *Science* **276**, 1221–1229 (1997).
- Zehr, J. P. & Paerl, H. W. in *Microbial Ecology of the Oceans* (ed. Kirchman, D. L.) 481–525 (John Wiley & Sons, New Jersey, 2008).
- Riemann, L., Farnelid, H. & Steward, G. F. Nitrogenase genes in non-cyanobacterial plankton: prevalence, diversity and regulation in marine waters. *Aquat. Microb. Ecol.* **61**, 235–247 (2010).
- Zehr, J. P. Nitrogen fixation by marine cyanobacteria. *Trends Microbiol.* **19**, 162–173 (2011).
- Berman-Frank, I. *et al.* Segregation of nitrogen fixation and oxygenic photosynthesis in the marine cyanobacterium *Trichodesmium*. *Science* **294**, 1534–1537 (2001).
- Breitbarth, E., Oschlies, A. & LaRoche, J. Physiological constraints on the global distribution of *Trichodesmium* — effect of temperature on diazotrophy. *Biogeosciences* **4**, 53–61 (2007).
- Zehr, J. P. *et al.* Unicellular cyanobacteria fix N₂ in the subtropical North Pacific Ocean. *Nature* **412**, 635–638 (2001).
- Church, M. J., Bjorkman, K. M., Karl, D. M., Saito, M. A. & Zehr, J. P. Regional distributions of nitrogen-fixing bacteria in the Pacific Ocean. *Limnol. Oceanogr.* **53**, 63–77 (2008).
- Langlois, R. J., Hummer, D. & LaRoche, J. Abundances and distributions of the dominant *nifH* phylotypes in the Northern Atlantic Ocean. *Appl. Environ. Microbiol.* **74**, 1922–1931 (2008).
- Goebel, N. L. *et al.* Abundance and distribution of major groups of diazotrophic cyanobacteria and their potential contribution to N₂ fixation in the tropical Atlantic Ocean. *Environ. Microbiol.* **12**, 3272–3289 (2010).

15. Montoya, J. P. *et al.* High rates of N₂ fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. *Nature* **430**, 1027–1031 (2004).
16. Webb, E. A., Ehrenreich, I. M., Brown, S. L., Valois, F. W. & Waterbury, J. B. Phenotypic and genotypic characterization of multiple strains of the diazotrophic cyanobacterium, *Crocosphaera watsonii*, isolated from the open ocean. *Environ. Microbiol.* **11**, 338–348 (2009).
17. Zehr, J. P., Bench, S. R., Mondragon, E. A., McCarren, J. & DeLong, E. F. Low genomic diversity in tropical oceanic N₂-fixing cyanobacteria. *Proc. Natl Acad. Sci. USA* **104**, 17807–17812 (2007).
18. Moisaner, P. H. *et al.* Unicellular cyanobacterial distributions broaden the oceanic N₂ fixation domain. *Science* **327**, 1512–1514 (2010).
19. Tripp, H. J. *et al.* Metabolic streamlining in an open-ocean nitrogen-fixing cyanobacterium. *Nature* **464**, 90–94 (2010).
- An excellent paper describing the genome of an important uncultured cyanobacterium, which was assembled from sorted environmental samples, and detailing its novel metabolism.**
20. Zehr, J. P. *et al.* Globally distributed uncultivated oceanic N₂-fixing cyanobacteria lack oxygenic Photosystem II. *Science* **322**, 1110–1112 (2008).
21. Needoba, J. A., Foster, R. A., Sakamoto, C., Zehr, J. P. & Johnson, K. S. Nitrogen fixation by unicellular diazotrophic cyanobacteria in the temperate oligotrophic North Pacific Ocean. *Limnol. Oceanogr.* **52**, 1317–1327 (2007).
22. Lehtimäki, J., Moisaner, P., Sivonen, K. & Kononen, K. Growth, nitrogen fixation, and nodularin production by two Baltic sea cyanobacteria. *Appl. Environ. Microbiol.* **63**, 1647–1656 (1997).
23. Laamanen, M., Kuosa, H. & Maximus, S. Annual variability of biomass and heterocysts in the N₂-fixing cyanobacterium *Aphanizomenon flos-aquae* in the Baltic Sea with reference to *Anabaena* spp. and *Nodularia spumigena*. *Boreal Environ. Res.* **10**, 19–30 (2005).
24. Foster, R. A. & Zehr, J. P. Characterization of diatom-cyanobacteria symbioses on the basis of *nifH*, *hetR* and 16S rRNA sequences. *Environ. Microbiol.* **8**, 1913–1925 (2006).
25. Bar Zeev, E. *et al.* Seasonal dynamics of the endosymbiotic, nitrogen-fixing cyanobacterium *Richelia intracellularis* in the eastern Mediterranean Sea. *ISME J.* **2**, 911–923 (2008).
26. Carpenter, E. J. *et al.* Extensive bloom of a N₂ fixing diatom/cyanobacterial association in the tropical Atlantic Ocean. *Mar. Ecol. Prog. Ser.* **185**, 273–283 (1999).
27. Venrick, E. L. Distribution and significance of *Richelia intracellularis* Schmidt in North Pacific central gyre. *Limnol. Oceanogr.* **19**, 437–445 (1974).
28. Subramaniam, A. *et al.* Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean. *Proc. Natl Acad. Sci. USA* **105**, 10460–10465 (2008).
- A paper detailing how the Amazon River enhances N₂ fixation and how the different diazotrophs present affect carbon sequestration in the area.**
29. Falkowski, P. G., Barber, R. T. & Smetacek, V. Biogeochemical controls and feedbacks on ocean primary production. *Science* **281**, 200–206 (1998).
30. Kustka, A. B. *et al.* Iron requirements for dinitrogen- and ammonium-supported growth in cultures of *Trichodesmium* (IMS 101): comparison with nitrogen fixation rates and iron: carbon ratios of field populations. *Limnol. Oceanogr.* **48**, 1869–1884 (2003).
31. Berman-Frank, I., Cullen, J. T., Shaked, Y., Sherrill, R. M. & Falkowski, P. G. Iron availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*. *Limnol. Oceanogr.* **46**, 1249–1260 (2001).
32. Bergquist, B. A. & Boyle, E. A. Dissolved iron in the tropical and subtropical Atlantic Ocean. *Global Biogeochem. Cycles* **20**, GB1015 (2006).
33. Blain, S., Bonnet, S. & Guieu, C. Dissolved iron distribution in the tropical and sub tropical South Eastern Pacific. *Biogeochemistry* **5**, 269–280 (2008).
34. Boyle, E. A., Bergquist, B. A., Kayser, R. A. & Mahowald, N. Iron, manganese, and lead at Hawaii Ocean Time-series station ALOHA: temporal variability and an intermediate water hydrothermal plume. *Geochim. Cosmochim. Acta* **69**, 933–952 (2005).
35. Wu, J. F., Boyle, E., Sunda, W. & Wen, L. S. Soluble and colloidal iron in the oligotrophic North Atlantic and North Pacific. *Science* **293**, 847–849 (2001).
36. Paerl, H. W., Prufertbebout, L. E. & Guo, C. Z. Iron-stimulated N₂ fixation and growth in natural and cultured populations of the planktonic marine cyanobacteria *Trichodesmium* spp. *Appl. Environ. Microbiol.* **60**, 1044–1047 (1994).
37. Rueter, J. G. Iron stimulation of photosynthesis and nitrogen-fixation in *Anabaena*-7120 and *Trichodesmium* (Cyanophyceae). *J. Phycol.* **24**, 249–254 (1988).
38. Chappell, P. D. & Webb, E. A. A molecular assessment of the iron stress response in the two phylogenetic clades of *Trichodesmium*. *Environ. Microbiol.* **12**, 13–27 (2010).
39. Kupper, H. *et al.* Iron limitation in the marine cyanobacterium *Trichodesmium* reveals new insights into regulation of photosynthesis and nitrogen fixation. *New Phytol.* **179**, 784–798 (2008).
40. Shi, T., Sun, Y. & Falkowski, P. G. Effects of iron limitation on the expression of metabolic genes in the marine cyanobacterium *Trichodesmium erythraeum* IMS101. *Environ. Microbiol.* **9**, 2945–2956 (2007).
41. Fu, F. X. *et al.* Interactions between pCO₂, N₂ fixation, and Fe limitation in the marine unicellular cyanobacterium *Crocosphaera*. *Limnol. Oceanogr.* **53**, 2472–2484 (2008).
42. Saito, M. A. *et al.* Iron conservation by reduction of metalloenzyme inventories in the marine diazotroph *Crocosphaera watsonii*. *Proc. Natl Acad. Sci. USA* **108**, 2184–2189 (2011).
43. Paczuska, L. & Kosakowska, A. Is iron a limiting factor of *Nodularia spumigena* blooms? *Oceanologia* **45**, 679–692 (2003).
44. Van Mooy, B. A. S. *et al.* Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. *Nature* **458**, 69–72 (2009).
45. Fu, F. X., Zhang, Y. H., Bell, P. R. F. & Hutchins, D. A. Phosphate uptake and growth kinetics of *Trichodesmium* (Cyanobacteria) isolates from the North Atlantic Ocean and the Great Barrier Reef, Australia. *J. Phycol.* **41**, 62–73 (2005).
46. Mulholland, M. R., Fløge, S., Carpenter, E. J. & Capone, D. G. Phosphorus dynamics in cultures and natural populations of *Trichodesmium* spp. *Mar. Ecol. Prog. Ser.* **239**, 45–55 (2002).
47. Krauk, J. M., Villareal, T. A., Sohm, J. A., Montoya, J. P. & Capone, D. G. Plasticity of N: P ratios in laboratory and field populations of *Trichodesmium* spp. *Aquat. Microb. Ecol.* **42**, 243–253 (2006).
48. White, A. E., Spitz, Y. H., Karl, D. M. & Letelier, R. M. Flexible elemental stoichiometry in *Trichodesmium* spp. and its ecological implications. *Limnol. Oceanogr.* **51**, 1777–1790 (2006).
49. Orchard, E. D., Webb, E. A. & Dyrhman, S. T. Molecular analysis of the phosphorus starvation response in *Trichodesmium* spp. *Environ. Microbiol.* **11**, 2400–2411 (2009).
50. Dyrhman, S. T. *et al.* Phosphonate utilization by the globally important marine diazotroph *Trichodesmium*. *Nature* **439**, 68–71 (2006).
51. Beversdorf, L. J., White, A. E., Bjorkman, K. M., Letelier, R. M. & Karl, D. M. Phosphonate metabolism by *Trichodesmium* IMS101 and the production of greenhouse gases. *Limnol. Oceanogr.* **55**, 1768–1778 (2010).
52. Dyrhman, S. T. & Haley, S. T. Phosphorus scavenging in the unicellular marine diazotroph *Crocosphaera watsonii*. *Appl. Environ. Microbiol.* **72**, 1452–1458 (2006).
53. Degerholm, J., Gundersen, K., Bergman, B. & Soderback, E. Phosphorus-limited growth dynamics in two Baltic Sea cyanobacteria, *Nodularia* sp and *Aphanizomenon* sp. *FEMS Microbiol. Ecol.* **58**, 323–332 (2006).
54. Mills, M. M., Ridame, C., Davey, M., La Roche, J. & Geider, R. J. Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic. *Nature* **429**, 292–294 (2004).
55. Moore, C. M. *et al.* Large-scale distribution of Atlantic nitrogen fixation controlled by iron availability. *Nature Geosci.* **2**, 867–871 (2009).
- An excellent paper synthesizing new data with that available from databases to detail the mechanisms that control N₂ fixation in the Atlantic Ocean.**
56. Cavender-Bares, K. K., Karl, D. M. & Chisholm, S. W. Nutrient gradients in the western North Atlantic Ocean: relationship to microbial community structure and comparison to patterns in the Pacific Ocean. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **48**, 2373–2395 (2001).
57. Wu, J. F., Sunda, W., Boyle, E. A. & Karl, D. M. Phosphate depletion in the western North Atlantic Ocean. *Science* **289**, 759–762 (2000).
58. Sohm, J. A., Mahaffey, C. & Capone, D. G. Assessment of relative phosphorus limitation of *Trichodesmium* spp. in the North Pacific, North Atlantic, and the north coast of Australia. *Limnol. Oceanogr.* **53**, 2495–2502 (2008).
59. Webb, E. A., Jakuba, R. W., Moffett, J. W. & Dyrhman, S. T. Molecular assessment of phosphorus and iron physiology in *Trichodesmium* populations from the western Central and western South Atlantic. *Limnol. Oceanogr.* **52**, 2221–2232 (2007).
60. Sanudo-Wilhelmy, S. A. *et al.* Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central Atlantic Ocean. *Nature* **411**, 66–69 (2001).
61. Sohm, J. A. & Capone, D. G. Zonal differences in phosphorus pools, turnover and deficiency across the tropical North Atlantic Ocean. *Global Biogeochem. Cycles* **24**, GB2008 (2010).
62. Brown, M. T., Landing, W. M. & Measures, C. I. Dissolved and particulate Fe in the western and central North Pacific: results from the 2002 IOC cruise. *Geochimistry Geophysics Geosystems* **6**, Q10001 (2005).
63. Mahowald, N. M. *et al.* Atmospheric global dust cycle and iron inputs to the ocean. *Global Biogeochem. Cycles* **19**, GB4025 (2005).
64. Grabowski, M. N. W., Church, M. J. & Karl, D. M. Nitrogen fixation rates and controls at Stn ALOHA. *Aquat. Microb. Ecol.* **52**, 175–183 (2008).
65. Kitajima, S., Furuya, K., Hashihama, F., Takeda, S. & Kanda, J. Latitudinal distribution of diazotrophs and their nitrogen fixation in the tropical and subtropical western North Pacific. *Limnol. Oceanogr.* **54**, 537–547 (2009).
66. Karl, D. M., Bidigare, R. R. & Letelier, R. M. Long-term changes in plankton community structure and productivity in the North Pacific Subtropical Gyre: the domain shift hypothesis. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **48**, 1449–1470 (2001).
67. Hynes, A. M., Chappell, P. D., Dyrhman, S. T., Doney, S. C. & Webb, E. A. Cross-basin comparison of phosphorus stress and nitrogen fixation in *Trichodesmium*. *Limnol. Oceanogr.* **54**, 1438–1448 (2009).
68. Van Mooy, B. A. S. & Devoil, A. H. Assessing nutrient limitation of *Prochlorococcus* in the North Pacific subtropical gyre by using an RNA capture method. *Limnol. Oceanogr.* **53**, 78–88 (2008).
69. Moutin, T. *et al.* Phosphate availability and the ultimate control of new nitrogen input by nitrogen fixation in the tropical Pacific Ocean. *Biogeochemistry* **5**, 95–109 (2008).
70. Raimbault, P. & Garcia, N. Evidence for efficient regenerated production and dinitrogen fixation in nitrogen-deficient waters of the South Pacific Ocean: impact on new and export production estimates. *Biogeochemistry* **5**, 323–338 (2008).
71. Stolte, W. *et al.* Stimulation of nitrogen-fixing cyanobacteria in a Baltic Sea plankton community by land-derived organic matter or iron addition. *Mar. Ecol. Prog. Ser.* **327**, 71–82 (2006).
72. Lilover, M. J. & Stips, A. The variability of parameters controlling the cyanobacteria bloom biomass in the Baltic Sea. *J. Mar. Sys.* **74**, S108–S115 (2008).
73. Moisaner, P. H., Paerl, H. W., Dyble, J. & Sivonen, K. Phosphorus limitation and diel control of nitrogen-fixing cyanobacteria in the Baltic Sea. *Mar. Ecol. Prog. Ser.* **345**, 41–50 (2007).
74. Nausch, M., Nausch, G. & Wasmund, N. Phosphorus dynamics during the transition from nitrogen to phosphate limitation in the central Baltic Sea. *Mar. Ecol. Prog. Ser.* **266**, 15–25 (2004).
75. Lips, I. & Lips, U. Abiotic factors influencing cyanobacterial bloom development in the Gulf of Finland (Baltic Sea). *Hydrobiologia* **614**, 133–140 (2008).
76. Nausch, M., Nausch, G., Wasmund, N. & Nagel, K. Phosphorus pool variations and their relation to cyanobacteria development in the Baltic Sea: a three-year study. *J. Mar. Sys.* **71**, 99–111 (2008).
77. Walve, J. & Larsson, U. Blooms of Baltic Sea *Aphanizomenon* sp (cyanobacteria) collapse after internal phosphorus depletion. *Aquat. Microb. Ecol.* **49**, 57–69 (2007).
78. Moisaner, P. H., Steppe, T. F., Hall, N. S., Kuparinen, J. & Paerl, H. W. Variability in nitrogen and phosphorus limitation for Baltic Sea phytoplankton during nitrogen-fixing cyanobacterial blooms. *Mar. Ecol. Prog. Ser.* **262**, 81–95 (2003).
79. Thingstad, T. F. *et al.* Nature of phosphorus limitation in the ultraoligotrophic eastern Mediterranean. *Science* **309**, 1068–1071 (2005).

80. Krom, M. D., Kress, N., Brenner, S. & Gordon, L. I. Phosphorus limitation of primary productivity in the eastern Mediterranean Sea. *Limnol. Oceanogr.* **36**, 424–432 (1991).
81. Man-Aharonovich, D., Kress, N., Bar Zeev, E., Berman-Frank, I. & Beja, O. Molecular ecology of *nifH* genes and transcripts in the eastern Mediterranean Sea. *Environ. Microbiol.* **9**, 2354–2363 (2007).
82. Le Moal, M. & Biegala, I. C. Diazotrophic unicellular cyanobacteria in the northwestern Mediterranean Sea: a seasonal cycle. *Limnol. Oceanogr.* **54**, 845–855 (2009).
83. Ibello, V., Cantoni, C., Cozzi, S. & Civitarese, G. First basin-wide experimental results on N₂ fixation in the open Mediterranean Sea. *Geophys. Res. Lett.* **37**, L03608 (2010).
84. Krom, M. D., Emeis, K. C. & Van Cappellen, P. Why is the Eastern Mediterranean phosphorus limited? *Prog. Oceanogr.* **85**, 236–244 (2010).
85. Mulholland, M. R. & Capone, D. G. In *Indian Ocean Biogeochemical Processes and Ecological Variability* (eds Wiggert, J. D., Hood, R. R., Naqvi, S. W. A., Brink, K. H. & Smith, S. L.) 167–186 (American Geophysical Union, Washington D. C., 2009).
86. Morrison, J. M. *et al.* Seasonal variation of hydrographic and nutrient fields during the US JGOFS Arabian Sea process study. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **45**, 2053–2101 (1998).
87. Capone, D. G. *et al.* An extensive bloom of the N₂ fixing cyanobacterium *Trichodesmium erythraeum* in the central Arabian Sea. *Mar. Ecol. Prog. Ser.* **172**, 281–292 (1998).
88. Foster, R. A., Paytan, A. & Zehr, J. P. Seasonality of N₂ fixation and *nifH* gene diversity in the Gulf of Aqaba (Red Sea). *Limnol. Oceanogr.* **54**, 219–233 (2009).
89. Carpenter, E. J., Subramaniam, A. & Capone, D. G. Biomass and primary productivity of the cyanobacterium *Trichodesmium* spp. in the tropical N Atlantic ocean. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **51**, 173–203 (2004).
90. Sohm, J. A., Subramaniam, A., Gunderson, T., Carpenter, E. J. & Capone, D. G. Nitrogen fixation by *Trichodesmium* spp. and unicellular diazotrophs in the North Pacific subtropical gyre. *J. Geophys. Res. Biogeosci.* 29 Mar 2011 (doi:10.1029/2010JG001513).
91. Bonnet, S., Biegala, I. C., Dutrieux, P., Slemmons, L. O. & Capone, D. G. Nitrogen fixation in the western equatorial Pacific: rates, diazotrophic cyanobacterial size class distribution, and biogeochemical significance. *Global Biogeochem. Cycles* **23**, GB3012 (2009).
92. Foster, R. A. *et al.* Influence of the Amazon River plume on distributions of free-living and symbiotic cyanobacteria in the western tropical north Atlantic Ocean. *Limnol. Oceanogr.* **52**, 2517–2532 (2007).
93. Foster, R. A., Subramaniam, A. & Zehr, J. P. Distribution and activity of diazotrophs in the Eastern Equatorial Atlantic. *Environ. Microbiol.* **11**, 741–750 (2009).
94. O’Neil, J. M. The colonial cyanobacterium *Trichodesmium* as a physical and nutritional substrate for the harpacticoid copepod *Macrosetella gracilis*. *J. Plankton Res.* **20**, 43–59 (1998).
95. Mulholland, M. R., Bronk, D. A. & Capone, D. G. Dinitrogen fixation and release of ammonium and dissolved organic nitrogen by *Trichodesmium* IMS101. *Aquat. Microb. Ecol.* **37**, 85–94 (2004).
96. Bordeleau, L. M. & Prevost, D. Nodulation and nitrogen fixation in extreme environments. *Plant Soil* **161**, 115–125 (1994).
97. Stal, L. J. Is the distribution of nitrogen-fixing cyanobacteria in the oceans related to temperature? *Environ. Microbiol.* **11**, 1632–1645 (2009). **An intriguing paper presenting a hypothesis to explain why free-living heterocystous cyanobacteria are found in cold waters and non-heterocystous filamentous cyanobacteria are found in warm waters.**
98. Holl, C. M., Waite, A. M., Pesant, S., Thompson, P. A. & Montoya, J. P. Unicellular diazotrophy as a source of nitrogen to Leeuwin Current coastal eddies. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **54**, 1045–1054 (2007).
99. Rees, A. P., Gilbert, J. A. & Kelly-Gerrey, B. A. Nitrogen fixation in the western English Channel (NE Atlantic Ocean). *Mar. Ecol. Prog. Ser.* **374**, 7–12 (2009).
100. Barcelos e Ramos, J., Biswas, H., Schulz, K. G., LaRoche, J. & Riebesell, U. Effect of rising atmospheric carbon dioxide on the marine nitrogen fixing *Trichodesmium*. *Global Biogeochem. Cycles* **21**, GB2028 (2007).
101. Hutchins, D. A. *et al.* CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth rates, and elemental ratios: implications for past, present, and future ocean biogeochemistry. *Limnol. Oceanogr.* **52**, 1293–1304 (2007).
102. Levitan, O. *et al.* Elevated CO₂ enhances nitrogen fixation and growth in the marine cyanobacterium *Trichodesmium*. *Glob. Change Biol.* **13**, 531–538 (2007).
103. Levitan, O. *et al.* Regulation of nitrogen metabolism in the marine diazotroph *Trichodesmium* IMS101 under varying temperatures and atmospheric CO₂ concentrations. *Environ. Microbiol.* **12**, 1899–1912 (2010).
104. Kranz, S. A. *et al.* Combined effects of CO₂ and light on the N₂ fixing Cyanobacterium *Trichodesmium* IMS101: physiological responses. *Plant Physiol.* **154**, 334–345 (2010).
105. Czerny, J., Ramos, J. B. E. & Riebesell, U. Influence of elevated CO₂ concentrations on cell division and nitrogen fixation rates in the bloom-forming cyanobacterium *Nodularia spumigena*. *Biogeosciences* **6**, 1865–1875 (2009).
106. Ramos, J. L., Madueno, F. & Guerrero, M. G. Regulation of nitrogenase levels in *Anabaena* Sp ATCC 3047 and other filamentous cyanobacteria. *Arch. Microbiol.* **141**, 105–111 (1985).
107. Martin-Nieto, J., Herrero, A. & Flores, E. Control of nitrogenase mRNA levels by products of nitrate assimilation in the cyanobacterium *Anabaena* sp. strain PCC-7120. *Plant Physiol.* **97**, 825–828 (1991).
108. Sanz-Alferez, S. & Del Campo, F. F. Relationship between nitrogen-fixation and nitrate metabolism in the *Nodularia* strains M1 and M2. *Planta* **194**, 339–345 (1994).
109. Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N. & Dunne, J. P. Spatial coupling of nitrogen inputs and losses in the ocean. *Nature* **445**, 163–167 (2007). **An exciting modelling effort presenting a new hypothesis on the location of the greatest N₂ fixation in the ocean, suggesting that N₂ fixation and denitrification are closely spatially coupled.**
110. Capone, D. G., Oneil, J. M., Zehr, J. & Carpenter, E. J. Basis for diel variation in nitrogenase activity in the marine planktonic cyanobacterium *Trichodesmium-thiebautii*. *Appl. Environ. Microbiol.* **56**, 3532–3536 (1990).
111. Fu, F. X. & Bell, P. R. F. Factors affecting N₂ fixation by the cyanobacterium *Trichodesmium* sp GBR-TRL101. *FEMS Microbiol. Ecol.* **45**, 203–209 (2003).
112. Mulholland, M. R., Ohki, K. & Capone, D. G. Nutrient controls on nitrogen uptake and metabolism by natural populations and cultures of *Trichodesmium* (Cyanobacteria). *J. Phycol.* **37**, 1001–1009 (2001).
113. Holl, C. M. & Montoya, J. P. Interactions between nitrate uptake and nitrogen fixation in continuous cultures of the marine diazotroph *Trichodesmium* (Cyanobacteria). *J. Phycol.* **41**, 1178–1183 (2005).
114. Davis, C. S. & McGillicuddy, D. J. Transatlantic abundance of the N₂ fixing colonial cyanobacterium *Trichodesmium*. *Science* **312**, 1517–1520 (2006).
115. Fong, A. A. *et al.* Nitrogen fixation in an anticyclonic eddy in the oligotrophic North Pacific Ocean. *ISME J.* **2**, 663–676 (2008).
116. Church, M. J. *et al.* Physical forcing of nitrogen fixation and diazotroph community structure in the North Pacific subtropical gyre. *Global Biogeochem. Cycles* **23**, GB2020 (2009).
117. Church, M. J., Short, C. M., Jenkins, B. D., Karl, D. M. & Zehr, J. P. Temporal patterns of nitrogenase gene (*nifH*) expression in the oligotrophic North Pacific Ocean. *Appl. Environ. Microbiol.* **71**, 5362–5370 (2005).
118. Moisander, P. H., Beinart, R. A., Voss, M. & Zehr, J. P. Diversity and abundance of diazotrophic microorganisms in the South China Sea during intermonsoon. *ISME J.* **2**, 954–967 (2008).
119. Moore, J. K., Doney, S. C. & Lindsay, K. Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model. *Global Biogeochem. Cycles* **18**, GB4028 (2004).
- A modelling study assessing nutrient limitation of different phytoplankton functional groups in the upper ocean.**
120. Duce, R. A. *et al.* Impacts of atmospheric anthropogenic nitrogen on the open ocean. *Science* **320**, 893–897 (2008).
121. Redfield, A. C. The biological control of chemical factors in the environment. *Am. Sci.* **46**, 205–221 (1958).
122. Capone, D. G. In *Nitrogen in the Marine Environment* (eds Carpenter, E. J. & Capone, D. G.) 105–137 (Springer, New York, 1983).
123. Howarth, R. W., Marino, R., Lane, J. & Cole, J. J. Nitrogen-fixation in fresh-water, estuarine, and marine ecosystems. 1. Rates and importance. *Limnol. Oceanogr.* **33**, 669–687 (1988).
124. Fulweiler, R. W., Nixon, S. W., Buckley, B. A. & Granger, S. L. Reversal of the net dinitrogen gas flux in coastal marine sediments. *Nature* **448**, 180–182 (2007).
125. Bertics, V. J. *et al.* Burrowing deeper into benthic nitrogen cycling: the impact of bioturbation on nitrogen fixation coupled to sulfate reduction. *Mar. Ecol. Prog. Ser.* **409**, 1–15 (2010).
126. Dekas, A. E., Poretsky, R. S. & Orphan, V. J. Deep-sea archaea fix and share nitrogen in methane-consuming microbial consortia. *Science* **326**, 422–426 (2009).
127. Mehta, M. P. & Baross, J. A. Nitrogen fixation at 92 °C by a hydrothermal vent archaeon. *Science* **314**, 1783–1786 (2006).
128. Severin, I. & Stal, L. J. *NifH* expression by five groups of phototrophs compared with nitrogenase activity in coastal microbial mats. *FEMS Microbiol. Ecol.* **73**, 55–67 (2010).
129. Moffett, J. W., Goeffert, T. J. & Naqvi, S. W. A. Reduced iron associated with secondary nitrite maxima in the Arabian Sea. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **54**, 1341–1349 (2007).
130. Bonnet, S. & Guieu, C. Atmospheric forcing on the annual iron cycle in the western Mediterranean Sea: a 1-year survey. *J. Geophys. Res.* **111**, C09010 (2006).
131. Breitbarth, E. *et al.* Dissolved iron (II) in the Baltic Sea surface water and implications for cyanobacterial bloom development. *Biogeosciences* **6**, 2397–2420 (2009).
132. Karl, D. M. *et al.* Ecological nitrogen-to-phosphorus stoichiometry at station ALOHA. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **48**, 1529–1566 (2001).
133. Dore, J. E., Brum, J. R., Tupas, L. M. & Karl, D. M. Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean. *Limnol. Oceanogr.* **47**, 1595–1607 (2002).
134. Wasmund, N., Voss, M. & Lochte, K. Evidence of nitrogen fixation by non-heterocystous cyanobacteria in the Baltic Sea and re-calculation of a budget of nitrogen fixation. *Mar. Ecol. Prog. Ser.* **214**, 1–14 (2001).
135. Tanaka, T., Rassoulzadegan, F. & Thingstad, T. F. Orthophosphate uptake by heterotrophic bacteria, cyanobacteria, and autotrophic nanoflagellates in Villefranche Bay, northwestern Mediterranean: vertical, seasonal, and short-term variations of the competitive relationship for phosphorus. *Limnol. Oceanogr.* **49**, 1063–1072 (2004).

Acknowledgements

The authors would like to express their thanks to M. Church, R. Foster, R. Langlois, J. LaRoche, P. Moisander and J. Zehr for providing *Trichodesmium* spp. *nifH* data and J. K. Moore for a model output, both of which were included in a previous version of this manuscript. Three anonymous reviewers helped to greatly improve the manuscript.

Competing interests statement

The authors declare no competing financial interests.

FURTHER INFORMATION

Jill A. Sohm's homepage:
<http://www.scf.usc.edu/~sohm/>
 Eric A. Webb's homepage:
<http://dornsife.usc.edu/labs/webb/home/index.cfm>
 Douglas G. Capone's homepage:
<http://www.usc.edu/dept/LAS/biosci/tricho/>

ALL LINKS ARE ACTIVE IN THE ONLINE PDF