

Moving Algebraic Cycles of Bounded Degree

Eric M. Friedlander* and H. Blaine Lawson**

The Chow Moving Lemma is a theorem which asserts that a given algebraic s -cycle on a smooth algebraic variety X can be moved within its rational equivalence class to intersect properly a given r -cycle on X provided that $r + s \geq \dim(X)$ (cf. [Chow], [S2]). In the past few years, there has been considerable interest in studying spaces of algebraic cycles rather than simply cycles modulo an equivalence relation. With this in mind, it is natural to ask whether one can move a given “bounded family” of s -cycles on the smooth variety X to intersect properly a given “bounded family” of r -cycles. The main point of this paper is to formulate and prove just such a result. In Theorem 3.1, we demonstrate that for any integer e and any smooth projective variety X , one can simultaneously and algebraically “move” all effective s -cycles of degree $\leq e$ on X so that each such cycle meets every effective r -cycle of degree $\leq e$ on X in proper dimension.

The primary motivation for this Moving Lemma for Cycles of Bounded Degree was the possibility of a duality theorem between cohomology and homology theories defined in terms of homotopy groups of cycle spaces. Using Theorem 3.1, we have proved such a duality theorem for complex quasi-projective varieties in [F-L2]. We prove our Moving Lemma for varieties over an arbitrary infinite field, permitting a proof in [F-V] of a duality theorem for “motivic cohomology and homology”.

The reader will find that our Moving Lemma has numerous good properties. First of all, the move is given as an algebraic move (parametrized by a punctured projective line) on Chow varieties. Although this move is “good” only for s -cycles of bounded degree, it is defined on all effective s -cycles. Moreover, the move starts at “time 0” by expressing an effective s -cycle Z as a difference of effective s -cycles both of which have intersection properties no worse than Z . Finally, our Moving Lemma is applicable to smooth quasi-projective varieties, for it is stated for a possibly singular projective variety X resulting in a conclusion of proper intersection off the singular locus of X .

The classical motivation for the moving lemma was to define an intersection product on algebraic cycles modulo rational equivalence, thereby establishing the Chow ring $A_*(X)$. Some of the classical literature overlooked the question of whether or not intersection of cycles defined via a moving lemma is independent of the move (e.g., [Chow], [R], [S2]; on the other hand, cf. [Chev], [S3]). One direct consequence of our Moving Lemma is a proof for smooth quasi-projective varieties that the intersection product is indeed well defined independent of the choice of move (Theorem 3.4). Of course, the intersection product now has an intrinsic formulation for all smooth algebraic varieties due to Fulton

*Partially supported by the N.S.F. and NSA Grant # MDA904-93-H-3029

**Partially supported by the N.S.F.

and MacPherson [Fu]. We should emphasize that our arguments are projective in nature, and thus apply directly only to quasi-projective varieties.

We gratefully acknowledge helpful comments from William Fulton, Andrei Suslin, and Vladimir Voevodsky. We are especially indebted to Ofer Gabber who foresaw the validity of such a moving lemma for families and directed us to local estimates arising in the proof of the key Theorem 1.1.

0. Definitions, Conventions, and a Brief Outline

We adopt the following conventions throughout. We fix a ground field k , choose an algebraic closure $\bar{k} \supset k$, and consider projective spaces \mathbf{P}^n over k of various dimensions n . A **projective variety** over k is a reduced and absolutely irreducible algebraic k -scheme which admits a (Zariski) closed embedding in some projective space \mathbf{P}^n . A **quasi-projective variety** over k is a Zariski open subset of some projective algebraic variety over k whose complement has defining ideal also defined over k . The set of **geometric points** $X(\bar{k})$ of such a quasi-projective variety X is the set of morphisms $\text{Spec}(\bar{k}) \rightarrow X$ of schemes over k . For a point x on a quasi-projective variety X , we denote by $\mathcal{O}_{X,x}$ the stalk at $x \in X$ of the structure sheaf of X and by \mathfrak{m}_x the maximal ideal of this local ring. An **algebraic cycle** Z of dimension r on a quasi-projective variety X is an integral combination of closed, r -dimensional subvarieties of X , which are (reduced, irreducible but not necessarily absolutely irreducible and) defined over k . We say Z is **effective** if the integer coefficients are all positive. Given a cycle Z on X , we define its **support** to be the algebraic subset $|Z| \subset X$ consisting of the union of the irreducible components of Z . If Y and Z are cycles on X of dimension r and s respectively with $r + s \geq m = \dim(X)$, we say that Y and Z **intersect properly** if each component of $|Y| \cap |Z|$ has dimension $\leq r + s - m$. If X is smooth and if Y and Z intersect properly, then the **intersection product** $Y \bullet Z$ is a well-defined cycle of dimension $r + s - m$ on X (See [Fu]).

Let $X \subset \mathbf{P}^n$ be a closed embedding over k of an m -dimensional projective variety X . For integers $d \geq 0$ and r with $0 \leq r \leq \dim(X)$, we denote by $\mathcal{C}_{r,d}(X)$ the **Chow variety** of effective r -cycles of degree d on X (cf. [S1]). The disjoint union

$$\mathcal{C}_r(X) = \coprod_{d \geq 0} \mathcal{C}_{r,d}(X)$$

has the structure of an abelian monoid and is called the **Chow monoid**. The geometric points $\mathcal{C}_r(X)(\bar{k})$ of this Chow monoid constitute the (discrete) monoid of effective r -cycles on $X_{\bar{k}}$, the base-change of X to $\text{Spec}(\bar{k})$. (See [F,1.2] for a discussion of the the k' -rational points of $\mathcal{C}_r(X)$ for any k'/k .)

Given an r -cycle Y and an s -cycle Z on a projective variety $X \subset \mathbf{P}^n$ of dimension m with $r + s \geq m$, the classical Chow Moving Lemma (cf. [R]) asserts the existence of a rational equivalence between Z and a necessarily ineffective s -cycle Z' which intersects Y properly at all smooth points of X . Roughly speaking the proof proceeds as follows. One considers a linear projection $\pi_L : \mathbf{P}^n \rightarrow \mathbf{P}^m$ whose vertex $L \cong \mathbf{P}^{n-m-1}$ does not meet X and constructs the **projecting cone** $C_L(Z) = \pi_L^*(\pi_{L*}(Z))$ of Z . Since $\pi_L|_X$ is a finite morphism, $C_L(X) \bullet X$ is a well-defined s -cycle on X which can be moved to intersect

Y properly by moving the cone $C_L(Z)$ in \mathbf{P}^n . The idea then is to choose L so that the **residual cycle**

$$R_L(Z) = C_L(Z) \bullet X - Z$$

intersects Y properly at least at all points outside the ramification locus of $p_L \stackrel{\text{def}}{=} \pi_L|_X$. Furthermore, any component of excess intersection of $|R_L(Z)| \cap |Y|$ lies in $\text{ram}(p_L) \cap |Z| \cap |Y|$. Thus, if we choose L_1, \dots, L_{m+1} so that $\cap_i \text{ram}(p_{L_i})$ is contained in the singular locus of X , then after $m + 1$ iterations the residual cycle $R_{L_{m+1}} \circ R_{L_m} \circ \dots \circ R_{L_1}(Z)$ will meet Y properly at all regular points of X .

In our paper we shall carry through this argument so that it applies simultaneously to all effective cycles Y, Z of degree bounded by any specified positive integer e . To do this we consider the reembeddings $X \hookrightarrow \mathbf{P}^{M_d}$ via the Veronese embeddings $\mathbf{P}^n \hookrightarrow \mathbf{P}^{M_d}$ of degree d . For each d we consider the subset $\mathcal{U}_X(d) \subset \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)^{m+1}))$ of those $(m + 1)$ -tuples of degree d hyperplanes whose intersection does not meet X . The main point is to show that the codimension of “bad L ’s” in $\mathcal{U}_X(d)$, i.e., those for which $R_L(Z)$ does not have an improved intersection with Y , goes to infinity as d goes to infinity. This is achieved in §1.

In §2 we construct the moving of the projecting cones. This is conventionally done by projective transformations of the ambient \mathbf{P}^n . Here we introduce a different method which is related to the movings introduced in [L] and algebraicized in [F] to prove the “algebraic suspension theorem” in Lawson homology. This argument essentially verifies that effective s -cycles of bounded degree in \mathbf{P}^n can be moved to intersect properly a hyperplane. Here we generalize this argument so that it applies to all effective r -cycles of bounded degree, where $s + r \geq n$, rather than to a single hyperplane.

In §3 we present our main results. In addition to Theorems 3.1 and 3.4 mentioned above, we prove in Theorem 3.5 that the intersection product on homotopy groups of cycle spaces on a smooth, projective complex variety (cf. [F-G]) can be represented by the intersection of families of cycles which have been moved to intersect properly. In Theorem 3.7, we present a basic ingredient of duality theorems for cycle spaces. It is the assertion that for a flat map $X \rightarrow B$ from a smooth variety, families of s -cycles on X with $s \geq \dim B$ can be moved to be equidimensional over B .

1. Residual cycles

Consider an m -dimensional projective variety X provided with a closed embedding $X \subset \mathbf{P}^n$ over k . Choose a positive integer d and define

$$\mathcal{U}_X \equiv \mathcal{U}_X(d) \subset \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)^{m+1}))$$

to be the Zariski open set of those $\mathbf{F} = (f_0, \dots, f_m)$ with the property that the zero locus $\mathcal{L}_{\mathbf{F}} = \{t \in \mathbf{P}^n : \mathbf{F}(t) = 0\}$ misses X and thus has codimension $m + 1$ in \mathbf{P}^n . (More precisely, \mathbf{F} is a k -rational point of the indicated Zariski open subset.) Each such \mathbf{F} determines a finite morphism (defined over k)

$$p_{\mathbf{F}} : X \longrightarrow \mathbf{P}^m \tag{1.0.1}$$

which can be viewed as follows. Let $v : \mathbf{P}^n \longrightarrow \mathbf{P}^M$ denote the Veronese embedding of degree d (so $M = \binom{n+d}{d} - 1$). Then each $\mathbf{F} \in \mathcal{U}_X$ determines a surjective linear projection

$$\pi_{\mathbf{F}} : \mathbf{P}^M \dashrightarrow \mathbf{P}^m \quad (1.0.2)$$

with the property that $p_{\mathbf{F}} = \pi_{\mathbf{F}} \circ v|_X$. The condition $\mathbf{F}(x) \neq 0$ for $x \in X$ is equivalent to the condition that the linear subspace $L(\mathbf{F}) \stackrel{\text{def}}{=} \text{the vertex of the projection } \pi_{\mathbf{F}}$, does not meet $v(X)$.

Suppose now that $Y, Z \subset X$ are closed algebraic subsets of pure dimension r and s respectively, where $r + s \geq m$. Let $Y \times Z - \Delta$ denote $(Y \times Z) \cap (X^2 - \text{diag}(X))$ and denote by $Y \star_{\mathbf{F}} Z$ the following closed subset of $Y \times Z - \Delta$:

$$Y \star_{\mathbf{F}} Z \equiv \{(y, z) : y \neq z, p_{\mathbf{F}}(y) = p_{\mathbf{F}}(z)\}$$

for any $\mathbf{F} \in \mathcal{U}_X$. We begin by investigating the condition on $\mathbf{F} \in \mathcal{U}_X$ that $Y \star_{\mathbf{F}} Z$ should have pure dimension $r + s - m$.

An interesting special case of the following theorem is the case in which Y equals Z , a cycle of dimension $r \geq m/2$ on the projective variety X of dimension m . Then, our theorem asserts for “most” projections of sufficiently high degree that the projection is injective on components of Z off a subset of dimension no greater than $2r - m$.

In the proof of Theorem 1.1, we use repeatedly the theorem of upper semi-continuity of dimensions of fibres of a map of algebraic varieties (cf. [Mu;I.8]).

Theorem 1.1 *Let $X \subset \mathbf{P}^n$ be a closed embedding over k of an m -dimensional projective variety X . Consider closed algebraic subsets $Y, Z \subset X$ of pure dimension r, s respectively, and assume that $r + s \geq m$. Then for any $\mathbf{F} \in \mathcal{U}_X(d)$, each irreducible component of $Y \star_{\mathbf{F}} Z$ has dimension $\geq r + s - m$.*

Fix any integer $N > 0$, and suppose for each closed point $w = (y, z) \in Y \times Z - \Delta$ and for each $\varphi \in \Gamma(\mathcal{O}_{\mathbf{P}^n}(d))$ with $\varphi(y) \neq 0 \neq \varphi(z)$ that the map

$$\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)) \longrightarrow \mathcal{O}_{Y,y}/\mathfrak{m}_y^{N+1} \times \mathcal{O}_{Z,z}/\mathfrak{m}_z^{N+1},$$

sending f to the restrictions of f/φ , is surjective. Then

$$\mathcal{B}(d)_{\{Y,Z\}} \subset \mathcal{U}_X(d)$$

has Zariski closure in $\mathcal{U}_X(d)$ of codimension at least $N - r - s$, where $\mathcal{B}(d)_{\{Y,Z\}}$ consists of those \mathbf{F} for which some component of $Y \star_{\mathbf{F}} Z$ has dimension $> r + s - m$.

Proof. For notational convenience, we write $\mathcal{B}_{\{Y,Z\}} \subset \mathcal{U}_X$ for $\mathcal{B}(d)_{\{Y,Z\}} \subset \mathcal{U}_X(d)$. Consider the subvariety (closed by upper semi-continuity)

$$\begin{aligned} \mathcal{B} &\equiv \{(y, z, \mathbf{F}) : \text{some component of } Y \star_{\mathbf{F}} Z \text{ has } \dim > r + s - m; (y, z) \in (Y \star_{\mathbf{F}} Z)\} \\ &\subset (Y \times Z - \Delta) \times \mathcal{U}_X, \end{aligned}$$

and let \mathcal{B}_w denote the fibre of $pr_{1,2} : \mathcal{B} \rightarrow (X \times Y) - \Delta$ above $w \in (X \times Y) - \Delta$. We shall prove that

$$\text{the codimension of } \mathcal{B}_w \text{ in } \{w\} \times \mathcal{U}_X \text{ is } \geq N \quad (1.1.1)$$

for all $w \in (X \times Y) - \Delta$. Since the image of \mathcal{B} under $pr_3 : \mathcal{B} \rightarrow \mathcal{U}_X$ equals $\mathcal{B}_{Y,Z}$ and since $\dim(Y \times Z) = r + s$, we conclude by applying the upper semi-continuity theorem to $pr_{1,2} : \mathcal{B} \rightarrow (X \times Y) - \Delta$ that the codimension of (the closure of) $\mathcal{B}_{Y,Z} \subset \mathcal{U}_X$ must be $\geq N - r - s$.

Fix $w = (y, z) \in (Y \times Z) - \Delta$ and observe that $\mathcal{B}_w \subset \mathcal{U}_X$ is the closed subset of those $\mathbf{F} = (f_0, \dots, f_m)$ for which some component of $Y \star_{\mathbf{F}} Z$ has dimension $> r + s - m$ at w (and such that for each point $x \in X$ there is some f_i with $f_i(x) \neq 0$). Observe that locally at w the variety $Y \star_{\mathbf{F}} Z$ is defined by the equations:

$$f_\ell(y')f_j(z') = f_j(y')f_\ell(z') \quad \text{for } 0 \leq \ell, j \leq m \quad (1.1.2)$$

in a neighborhood of $w = (y, z)$ in $Y \times Z$.

Let Γ denote $\Gamma(\mathcal{O}_{\mathbf{P}^n}(d))$ and let $\Gamma' \subset \Gamma$ denote the open subset of those f with $f(y) \neq 0 \neq f(z)$. Observe that for any $\mathbf{F} = (f_0, \dots, f_m) \in \mathcal{U}_X$ there is some pair ℓ, j with $0 \leq \ell < j \leq m$ and $(f_\ell + f_j)(y) \neq 0 \neq (f_\ell + f_j)(z)$. Thus, $\mathcal{U}_X \subset \mathbf{P}(\Gamma^{m+1})$ is contained in the union of images of finitely many translates by elements of $GL(\Gamma^{m+1})$ of $\Gamma' \times \Gamma^m$. Let $\mathcal{B}_{w,m} \subset \Gamma' \times \Gamma^m$ denote the closed subvariety of those \mathbf{F} for which the subvariety determined by (1.1.2) has some component of dimension $> r + s - m$ at w . Then the union of the images of these finitely many translates of $\mathcal{B}_{w,m}$ intersected with \mathcal{U}_X equals \mathcal{B}_w . Consequently, (1.1.2) is implied by the assertion

$$\mathcal{B}_{w,m} \subset \Gamma' \times \Gamma^m \text{ has codimension } \geq N. \quad (1.1.3)$$

(The codimension does not drop under the projection $\Gamma^{m+1} - \{0\} \rightarrow \mathbf{P}(\Gamma^{m+1})$ since $\mathcal{B}_{w,m}$ is a cone.) By the upper semi-continuity theorem once again, (1.1.3) will follow if we show for each $\varphi \in \Gamma'$ that

$$\mathcal{B}_{w,m,\varphi} \subset \{\varphi\} \times \Gamma^m \text{ has codimension } \geq N, \quad (1.1.4)$$

where $\mathcal{B}_{w,m,\varphi} \equiv \mathcal{B}_{w,m} \cap (\{\varphi\} \times \Gamma^m)$.

We proceed to prove (1.1.4) by induction. Consider for each i with $1 \leq i \leq m$ the set of equations

$$f_\ell(y')f_j(z') = f_j(y')f_\ell(z') \quad \text{for } 0 \leq \ell, j \leq i. \quad (1.1.5)_i$$

For $\mathbf{F} = (\varphi, f_1, \dots, f_m) \in \mathcal{B}_{w,i,\varphi}$, the set of equations (1.1.5)_i are equivalent at w to the set of equations:

$$g_{\mathbf{F}}^j(y', z') = \frac{f_j(y')}{\varphi(y')} - \frac{f_j(z')}{\varphi(z')} = 0 \quad \text{for } 1 \leq j \leq i. \quad (1.1.6)_i$$

In particular, we conclude that $Y \star_{\mathbf{F}} Z$ has dimension $\geq r + s - m$ in a neighborhood of w , since this is given by the m equations of (1.1.6)_m. We define $\mathcal{B}_{w,i,\varphi} \subset \mathcal{B}_{w,m,\varphi}$ to be

the subvariety of those $\mathbf{F} = (\varphi, f_1, \dots, f_m)$ such that some component of the subvariety of $X \times Y$ defined by (1.1.6)_{*i*} has dimension $> r + s - i$ at w . Further, we define $\mathcal{B}'_{w,i,\varphi} = \mathcal{B}_{w,i,\varphi} - \mathcal{B}_{w,i-1,\varphi}$. Using induction, we conclude that it suffices to prove

$$\mathcal{B}'_{w,i,\varphi} \subset \{\varphi\} \times \Gamma^m \text{ has codimension } \geq N. \quad (1.1.7)_i$$

To bound the codimension of $\mathcal{B}'_{w,i,\varphi}$, we rephrase the problem in terms of local rings. For any $\mathbf{F} \in \mathcal{B}_{w,i,\varphi}$, set $R(w, \mathbf{F})_0 = \mathcal{O}_{Y \times Z, w}$, and let $R(w, \mathbf{F})_i$ denote the local ring $\mathcal{O}_{Y \times Z, w} / \langle g_{\mathbf{F}}^1, \dots, g_{\mathbf{F}}^i \rangle$ for $1 \leq i \leq m$. If $R(w, \mathbf{F})_i$ has pure dimension $r + s - i$ for some $i < m$, then the condition on $g_{\mathbf{F}}^{i+1}$ that $R(w, \mathbf{F})_{i+1}$ have dimension $\leq r + s - i - 1$ is the condition that $g_{\mathbf{F}}^{i+1}$ be non-zero in $R(w, \mathbf{F})_{i,\alpha} \equiv R(w, \mathbf{F})_i / \mathcal{P}_{i,\alpha}$ for each of the finitely many minimal primes $\mathcal{P}_{i,\alpha}$ of $R(w, \mathbf{F})_i$. This is guaranteed if the class of $g_{\mathbf{F}}^{i+1}$ is non-zero in the quotient $R(w, \mathbf{F})_{i,\alpha} / \mathfrak{m}^{N+1}$ for all α , where \mathfrak{m} denotes the maximal ideal of $R(w, \mathbf{F})_{i,\alpha}$.

If $\mathbf{F} = (\varphi, f_1, \dots, f_m) \in \mathcal{B}'_{w,i+1,\varphi}$, then \mathbf{F} lies in the kernel of the linear map

$$G_{\alpha}^{i+1} : \{\varphi\} \times \Gamma^m \rightarrow R(w, \mathbf{F})_{i,\alpha} / \mathfrak{m}^{N+1}$$

sending \mathbf{F} to the image of $g_{\mathbf{F}}^{i+1}$. To prove (1.1.7)_{*i+1*} assuming (1.1.7)_{*i*}, it suffices to prove that $\text{rank}(G_{\alpha}^{i+1}) \geq N$.

Extend $R(w, \mathbf{F})_{i,\alpha}$ to a discrete valuation ring $R(w, \mathbf{F})_{i,\alpha} \subset A$ of its function field and let v_A denote the associated valuation. Let v denote $\min\{v_A(x) : x \in \mathfrak{m}\}$. Consider the composition

$$\mathcal{O}_{Y,y} / \mathfrak{m}_y^{N+1} \times \mathcal{O}_{Z,z} / \mathfrak{m}_z^{N+1} \rightarrow \mathcal{O}_{X \times Y, w} / \mathfrak{m}_w^{N+1} \rightarrow R(w, \mathbf{F})_{i,\alpha} / \mathfrak{m}^{N+1} \rightarrow A / \mathfrak{m}_A^{v(N+1)} \quad (1.1.8)$$

where the first map is difference map sending (f, g) to $f - g$. Since the images of the maximal ideals $\mathfrak{m}_y, \mathfrak{m}_z$ of $\mathcal{O}_{Y,y}, \mathcal{O}_{Z,z}$ generate the maximal ideal \mathfrak{m}_w of $\mathcal{O}_{X \times Y, w}$ and thus also \mathfrak{m} of $R(w, \mathbf{F})_{i,\alpha}$, we conclude that this composition has rank at least N . Our hypothesis on $\Gamma \rightarrow \mathcal{O}_{Y,y} / \mathfrak{m}_y^{N+1} \times \mathcal{O}_{Z,z} / \mathfrak{m}_z^{N+1}$ (sending a form $f \in \Gamma$ to the restrictions of f/φ) thus implies that G_{α}^{i+1} , defined as the composition of $\Gamma \rightarrow \mathcal{O}_{Y,y} / \mathfrak{m}_y^{N+1} \times \mathcal{O}_{Z,z} / \mathfrak{m}_z^{N+1}$ with (1.1.8), also has rank at least N . \square

In order to apply Theorem 1.1, we show in the following lemma that the surjectivity hypothesis of the proposition is satisfied for all sufficiently large degrees d .

Lemma 1.2. *Retain the notation of Theorem 1.1. For each $N > 0$, there exists some d_N such that for all $d \geq d_N$ the map*

$$\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)) \rightarrow \mathcal{O}_{Y,y} / \mathfrak{m}_y^{N+1} \times \mathcal{O}_{Z,z} / \mathfrak{m}_z^{N+1} \quad (1.2.1)$$

sending f to the restrictions of f/φ is surjective for all closed points $(y, z) \in Y \times Z - \Delta$, all $\varphi \in \Gamma(\mathcal{O}_{\mathbf{P}^n}(d))$ with $\varphi(y) \neq 0 \neq \varphi(z)$.

Proof. Since $\mathcal{O}_{\mathbf{P}^n, y} \rightarrow \mathcal{O}_{Y,y}, \mathcal{O}_{\mathbf{P}^n, z} \rightarrow \mathcal{O}_{Z,z}$ are surjective local homomorphisms, it suffices to consider the special case $Y = \mathbf{P}^n = Z$. Moreover, base-changing by a field extension $\text{Spec}(k') \rightarrow \text{Spec}(k)$ merely tensors the map (1.2.1) with k' over k . Consequently, it suffices

to assume that both y, z are rational points of \mathbf{P}^n . By the two-point homogeneity of \mathbf{P}^n under PGL_{n+1} , it thus suffices to verify the existence of d_N for a single pair of (rational) points $(y, z) \in \mathbf{P}^n \times \mathbf{P}^n$.

The map sending f to the restrictions of f/φ in $\mathcal{O}_{Y,y}/\mathfrak{m}_y^{N+1}$ and $\mathcal{O}_{Z,z}/\mathfrak{m}_z^{N+1}$ has image whose dimension is independent of the choice of φ with $\varphi(y) \neq 0 \neq \varphi(z)$ since the quotient of any two such choices of φ is invertible in both $\mathcal{O}_{Y,y}$ and $\mathcal{O}_{Z,z}$. Hence, it will suffice to consider the case where $\varphi = X_0^d$ for some $X_0 \in \Gamma(\mathcal{O}_{\mathbf{P}^n}(1))$.

Let $R = k[X_1/X_0, \dots, X_n/X_0]$ denote the coordinate algebra of the affine variety $\mathbf{A}^n = \mathbf{P}^n - \{X_0 = 0\}$, where we assume that $X_0(y) \neq 0 \neq X_0(z)$. Since $\mathfrak{m}_y^{N+1}, \mathfrak{m}_z^{N+1}$ are coprime in R , the Chinese Remainder Theorem implies that

$$R \longrightarrow R/\mathfrak{m}_y^{N+1} \times R/\mathfrak{m}_z^{N+1}$$

is surjective. Let d_N be the minimal degree d for which polynomials of degree $\leq d_N$ in R map via this surjective map onto $R/\mathfrak{m}_y^{N+1} \times R/\mathfrak{m}_z^{N+1}$. Then for any $d \geq d_N$, (1.2.1) is surjective. \square

Let X be a projective variety provided with a closed embedding $X \subset \mathbf{P}^n$ over k and consider non-negative integers r, d . For any non-negative integer e , set

$$\mathcal{C}_{r, \leq e}(X) = \coprod_{d \leq e} \mathcal{C}_{r,d}(X).$$

In the next proposition we consider the Veronese embeddings of X of degree d . We consider all cycles of fixed degree $\leq e$ (in the original embedding) and examine those projections which are “bad” for pairs of such cycles. The main assertion is that the codimension of these bad projections goes to infinity with d .

Proposition 1.3. *Let $X \subset \mathbf{P}^n$ be a closed subvariety of dimension m , let r, s be non-negative integers with $r + s \geq m$, and let e be a positive integer. There exist (Zariski) closed subsets $\mathcal{B}(d)_e$ of the quasi-projective variety $\mathcal{U}_X(d)$,*

$$\mathcal{B}(d)_e \subset \mathcal{U}_X(d) \subset \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)))^{m+1},$$

for $d > 0$ with

$$\lim_{d \rightarrow \infty} \text{codim} \mathcal{B}(d)_e = \infty$$

which satisfy the following property: for any effective r -cycle Y on X of degree $\leq e$ and any effective s -cycle Z on X also of degree $\leq e$, $|Y| \star_{\mathbf{F}} |Z|$ has pure dimension $r + s - m$ whenever \mathbf{F} is a k -rational point of $\mathcal{U}_X(d) - \mathcal{B}(d)_e$.

Proof. For notational brevity, let $\mathcal{C}_{r,e}$ denote $\mathcal{C}_{r, \leq e}(X)$. We define

$$\mathcal{W}_e \subset X^2 \times \mathcal{C}_{r,e} \times \mathcal{C}_{s,e} \times \mathcal{U}_X(d)$$

to be the closed subvariety of those quintuples (y, z, Y, Z, \mathbf{F}) for which $y \times z$ lies in the closure of some component of $Y \star_{\mathbf{F}} Z$. The principle of upper semi-continuity applied to \mathcal{W}_e over $\mathcal{C}_{r,e} \times \mathcal{C}_{s,e} \times \mathcal{U}_X(d)$ implies that

$$\mathcal{V}_e \subset \mathcal{C}_{r,e} \times \mathcal{C}_{s,e} \times \mathcal{U}_X(d)$$

is a closed subvariety, where \mathcal{V}_e consists of those triples (Y, Z, F) such that $Y \star_{\mathbf{F}} Z$ has some irreducible component of dimension $> r + s - m$. Thus,

$$\mathcal{B}(d)_e = \text{pr}_3(\mathcal{V}_e) \subset \mathcal{U}_X(d)$$

is a closed subvariety of $\mathcal{U}_X(d)$.

Observe that the fibre of \mathcal{V}_e above $(Y, Z) \in \mathcal{C}_r \times \mathcal{C}_s$ is $\mathcal{B}_{\{Y,Z\}}$ (in the notation of Theorem 1.1). Let $E_r = \dim\{\mathcal{C}_{r,e}\}, E_s = \dim\{\mathcal{C}_{s,e}\}$. By Theorem 1.1 and Lemma 1.2, for any N we may find d_N so that $\mathcal{B}(d)_{\{Y,Z\}} \subset \mathcal{U}_X(d)$ has codimension at least $N - E_r - E_s - r - s$ for $d \geq d_N$. Hence, $\mathcal{B}_e(d)$ has codimension $\geq N - r - s$ for $d \geq d_N$.

Finally, if Y, Z are effective r, s cycles of degrees $\leq e$ and if $Y \star_{\mathbf{F}} Z$ has some component of dimension $> r + s - m$, then (Y, Z, \mathbf{F}) determines a point of \mathcal{V}_e so that $\mathbf{F} \in \mathcal{B}(d)_e$. \square

Chow's classical technique of moving Z to intersect Y properly entails consideration of a Veronese embedding $\mathbf{P}^n \rightarrow \mathbf{P}^M$ and a linear plane $L \subset \mathbf{P}^M$ of codimension $m + 1$ missing X . We let $\pi_L : \mathbf{P}^M \dashrightarrow \mathbf{P}^m$ denote the linear projection with center L and $p_L : X \rightarrow \mathbf{P}^m$ denote the finite morphism given as the restriction of π_L to X . Departing from the classical construction (cf [R]), we define the **projecting cone** of a cycle Z on X to be the cycle

$$C_L(Z) \stackrel{\text{def}}{=} \pi_L^*(p_{L*}(Z)) \subset \mathbf{P}^M,$$

where p_{L*} denotes proper push-forward and π_L^* denotes the algebraic join operation $(-)\#L$ sending a cycle W on \mathbf{P}^m to $W\#L$ on $\mathbf{P}^m\#L = \mathbf{P}^M$ (cf. [L,2.10]). This differs from the classical definition which is given by the discontinuous construction of sending an irreducible cycle Z to $\pi_L^*(p_L(Z))$, where $p_L(Z)$ is the irreducible, reduced image of Z .

Since p_L is a finite map, the intersection product $C_L(Z) \bullet X$ is defined for all cycles Z on X . We recall that $p_L : X \rightarrow \mathbf{P}^m$ is flat at any smooth point $x \in X$ (cf. [Mat,20.D]). Thus, if no component of Z is contained in the singular locus of X , then $C_L(Z) \bullet X$ can alternatively be described as the closure in X of $p_L^{ns*}(p_{L*}(Z))$, where $p_L^{ns} : X_{ns} \rightarrow \mathbf{P}^m$ is the restriction to p_L to the complement of the singular locus of X .

One considers the **residual cycle**

$$R_L(Z) \stackrel{\text{def}}{=} C_L(Z) \bullet X - Z.$$

This is a linear (i.e., additive) construction on cycles, sending effective cycles to effective cycles (cf. Lemma 1.6).

Given Y and Z , one wants to choose L so that $R_L(Z)$ has an improved intersection with Y . A key to this is the following lemma proved in [R] for the classical definition of residual cycle involving the classical projecting cone $\pi_L^*(p_L(Z)) \bullet X$. Denote by $\text{ram}(p_L)$

the **ramification locus** of the finite map $p_L : X \longrightarrow \mathbf{P}^m$. (In particular $\text{ram}(p_L)$ contains all the singular points of X .)

Proposition 1.4 (cf. [R; Lemma 6]) *Let Z , X and L be as above and consider a geometric point $\underline{y} : \text{Spec}(\bar{k}) \rightarrow |R_L(Z)|$ above a closed point $y \in |R_L(Z)|$. Then either:*

- (i) *There exists a geometric point $\underline{z} : \text{Spec}(\bar{k}) \rightarrow |Z|$ over a closed point $z \in |Z|$ with $\underline{z} \neq \underline{y}$ and $p_L(\underline{z}) = p_L(\underline{y})$, or:*
- (ii) *$y \in \text{ram}(p_L)$.*

Proof. Since the construction of the residual cycle is additive, we may assume that Z is irreducible. By definition of $R_L(Z)$, if $y \notin Z$, then there exists a point $z \in Z$ with $z \neq y$ and $p_L(z) = p_L(y)$. Thus, we may assume $y \in Z$.

If $[Z : p_L(Z)] > 1$ (i.e., if the function field of Z is a non-trivial extension of that of $p_L(Z)$), then either (i) is satisfied for some $\underline{z} : \text{Spec}(\bar{k}) \rightarrow Z$ or in a formal neighborhood of \underline{y} at least two sheets of Z (above a formal neighborhood of $p_L(y) \in \mathbf{P}^m$) intersect at y . This latter situation implies that in a formal neighborhood of $y \in X$ at least two sheets of X intersect at y so that $y \in \text{ram}(p_L)$.

If $[Z : p_L(Z)] = 1$ and if Z is a component of $R_L(Z)$, then as shown in [R; Lemma 6] p_L must ramify along all of Z . Namely, a “general” point of Z is necessarily smooth on Z and would also be smooth on X if p_L does not ramify along Z . At such a point, p_L is etale and so the multiplicity of Z in $p_L^{n_s*}(p_{L*}(Z))$ is 1; this contradicts the assumption that Z is a component of $R_L(Z)$.

Finally, assume that $[Z : p_L(Z)] = 1$ and that Z is not a component of $R_L(Z)$. Since $y \in |R_L(Z)|$, y lies in some component R of $R_L(Z)$ not equal to Z . Then, in a formal neighborhood of $y \in |R_L(Z)|$ at least two sheets (one determined by Z and another determined by R) intersect at y so that $y \in \text{ram}(p_L)$. \square

As in (1.0.1) and (1.0.2), for each $(m+1)$ -tuple $\mathbf{F} = (f_0, \dots, f_m)$ of homogeneous forms of degree d on \mathbf{P}^n , we consider the associated linear projection $\pi_{\mathbf{F}} : \mathbf{P}^M \dashrightarrow \mathbf{P}^m$ with center $L(\mathbf{F})$. If $X \subset \mathbf{P}^n$ is a subvariety of dimension m and if $\mathbf{F} \in \mathcal{U}_X(d)$, we denote by $p_{\mathbf{F}} : X \rightarrow \mathbf{P}^m$ the finite map given by restricting $\pi_{\mathbf{F}}$ to $X \subset \mathbf{P}^M$ (embedded via the Veronese embedding). For any cycle Z on X we set

$$C_{\mathbf{F}}(Z) = \pi_{\mathbf{F}}^*(p_{\mathbf{F}*}((Z))) = C_{L(\mathbf{F})}(Z) \quad \text{and} \quad R_{\mathbf{F}}(Z) = C_{\mathbf{F}}(Z) \bullet X - Z.$$

Corollary 1.5. *Consider a projective variety X of dimension m provided with a closed embedding $X \subset \mathbf{P}^n$. Let Y be an r -cycle and Z an s -cycle on X with $r + s - m \geq 0$. Assume $\mathbf{F} \in \mathcal{U}_X(d)$ satisfies the condition that $|Y| \star_{\mathbf{F}} |Z|$ has no component of excess dimension (i.e., $> r + s - m$). Then any component of $|Y| \cap |R_{\mathbf{F}}(Z)|$ of excess dimension must be contained in $|Y| \cap |Z| \cap \text{ram}(p_{\mathbf{F}})$.*

Proof. By definition of $|Y| \star_{\mathbf{F}} |Z|$, any $y \in (|Y| \cap |R_{\mathbf{F}}(Z)|) - |Z|$ necessarily lies in $\text{pr}_{|Y|}(|Y| \star_{\mathbf{F}} |Z|)$. By Proposition 1.5, any $y \in (|Y| \cap |R_{\mathbf{F}}(Z)|) - \text{ram}(p_{\mathbf{F}})$ admits a geometric point $(\underline{y}, \underline{z}) : \text{Spec}(\bar{k}) \rightarrow |Y| \star_{\mathbf{F}} |Z|$ which projects to \underline{y} over y , so that y also lies in $\text{pr}_{|Y|}(|Y| \star_{\mathbf{F}} |Z|)$. Thus, the complement of $|Y| \cap |Z| \cap \text{ram}(p_{\mathbf{F}})$ in $|Y| \cap |R_{\mathbf{F}}(Z)|$ lies in

$\text{pr}_{|Y|}(|Y| \star_{\mathbf{F}} |Z|)$ whose components have dimensions less than or equal to that of the maximal dimension of the components of $|Y| \star_{\mathbf{F}} |Z|$. \square

We shall use the following property of the construction of residual cycles. We recall that a **continuous algebraic map** $f : X \rightarrow Y$ with Y projective is a set-theoretic map on geometric points which is induced by a correspondence $\Gamma_f \subset X \times Y$ (cf. [F]).

Lemma 1.6. *Let \mathbf{F} be a k -rational point of $\mathcal{U}_X(d) \subset \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^n}(d)^{m+1}))$. Then $R_{\mathbf{F}}(Z)$ is an effective s -cycle on X whenever Z is itself an effective s -cycle on X . Moreover, sending Z to $R_{\mathbf{F}}(Z)$ determines a continuous algebraic map*

$$R_{\mathbf{F}}(-) : \mathcal{C}_{s, \leq e}(X) \rightarrow \mathcal{C}_{s, \leq e'}(X),$$

where e' is a positive integer depending upon d, e and the degree of X .

Proof. The fact that $R_{\mathbf{F}}(-)$ sends effective cycles on X to effective cycles on X follows immediately from the observation that each of the operations in the definition of $R_{\mathbf{F}}(-)$ is a well defined operation on effective cycles.

We employ various functoriality properties proved in [F]. The Veronese embedding $\nu : \mathbf{P}^n \rightarrow \mathbf{P}^M$ of degree d determines a morphism

$$\nu_* : \mathcal{C}_{s, \leq e}(X) \rightarrow \mathcal{C}_{s, \leq d^s e}(\nu(X)).$$

The linear projection $p_{\mathbf{F}} : \nu(X) \rightarrow \mathbf{P}^m$ determines a morphism

$$p_{\mathbf{F}*} : \mathcal{C}_{s, \leq d^s e}(\nu(X)) \rightarrow \mathcal{C}_{s, \leq d^s e}(\mathbf{P}^m).$$

The join construction determines a continuous algebraic map

$$\pi_{\mathbf{F}}^* : \mathcal{C}_{s, \leq d^s e}(\mathbf{P}^m) \rightarrow \mathcal{C}_{s+M-m, \leq d^s e}(\mathbf{P}^M).$$

Furthermore, intersection product in \mathbf{P}^M determines a continuous algebraic map

$$(-) \bullet X : \mathcal{C}_{s+M-m, \leq d^s e}(\mathbf{P}^M) \rightarrow \mathcal{C}_{s, \leq d^s e f}(\mathbf{P}^M)$$

where f equals the degree of $\nu(X)$. Finally, the additive structure of the Chow monoid is also given by continuous algebraic maps. Thus, sending Z to $R_{\mathbf{F}}(Z) \equiv \pi_{\mathbf{F}}^*(p_{\mathbf{F}*}(Z)) \bullet X - Z$ is a continuous algebraic map. \square

In the following theorem, we verify that an iteration of the residual cycle construction enables one to arrange (by choosing the degrees of projections sufficiently large) that the iterated residual cycle for Z meets Y properly for all effective cycles Y, Z of degree $\leq e$.

Theorem 1.7 *Let $X \subset \mathbf{P}^n$ be a closed subvariety of dimension m , and let $U \subset X$ be the Zariski open subset consisting of the smooth points of X . For any $(m+1)$ -tuple of positive integers $\mathbf{d} = (d_0, \dots, d_m)$, there is a (Zariski) open dense subset*

$$\mathfrak{R}_X(\mathbf{d}) \subset \prod_{i=0}^m \mathcal{U}_X(d_i) \subset \prod_{i=0}^m \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^n}(d_i))^{m+1})$$

consisting of $(m + 1)$ -tuples $\mathbf{F}^* \equiv (\mathbf{F}^0, \dots, \mathbf{F}^m)$ with the property that

$$U \cap \text{ram}(p_{\mathbf{F}^0}) \cap \dots \cap \text{ram}(p_{\mathbf{F}^m}) = \emptyset.$$

For any non-negative integers r, s with $r + s \geq m$ and any positive integer e , there exists a (Zariski) closed subset

$$\mathfrak{B}(\mathbf{d})_e \subset \mathfrak{R}_X(\mathbf{d})$$

such that

- (a) The codimension of $\mathfrak{B}(\mathbf{d})_e$ can be made arbitrarily large for all appropriately chosen \mathbf{d} which are sufficiently large (cf. (1.8) below).
- (b) For all effective r -cycles Y and s -cycles Z of degree $\leq e$ on X and for all k -rational points $\mathbf{F}^* \in \mathfrak{R}_X(\mathbf{d}) - \mathfrak{B}(\mathbf{d})_e$, each component of

$$|Y| \cap |R_{\mathbf{F}^*}(Z)| \quad , \quad R_{\mathbf{F}^*}(Z) \equiv R_{\mathbf{F}^m} \circ \dots \circ R_{\mathbf{F}^0}(Z)$$

of “excess” dimension (i.e., $> r + s - m$) is contained in the singular locus $X - U$.

Proof. We consider the (closed) incidence correspondence $\mathcal{J} \subset \mathcal{U}_X(d_0) \times X \dots \times \mathcal{U}_X(d_m) \times X$ of those $(\mathbf{F}^0, x_0, \dots, \mathbf{F}^m, x_m)$ with $x_j \in \text{ram}(p_{\mathbf{F}^j})$ for all j . Then the complement of $\mathcal{R}_X(\mathbf{d})$ in $\mathcal{U}_X(\mathbf{d}) \equiv \prod_{j=0}^m \mathcal{U}_X(d_j)$ is given by

$$\mathcal{U}_X(\mathbf{d}) - \mathcal{R}_X(\mathbf{d}) = \text{pr}\{\mathcal{J} \cap (\mathcal{U}_X(\mathbf{d}) \times \Delta(X))\},$$

where pr is the proper projection $\text{pr} : \mathcal{U}_X(d_0) \times X \dots \times \mathcal{U}_X(d_m) \times X \rightarrow \mathcal{U}_X(\mathbf{d})$ and $\Delta : X \rightarrow X^{m+1}$ is the diagonal embedding. Hence, $\mathcal{R}_X(\mathbf{d})$ is open.

For any smooth point of $x \in X$, those $\mathbf{F} \in \mathcal{U}_X(d)$ such that $p_{\mathbf{F}}$ ramifies at x is a proper closed subset (defined by the condition that $L_{\mathbf{F}}$ meet the tangent plane of X at x). Thus, the subset of those \mathbf{F} for which $p_{\mathbf{F}}$ ramifies everywhere is a proper closed subset of $\mathcal{U}_X(d)$ (empty, if k has characteristic 0). We readily see that $\mathcal{R}_X(\mathbf{d})$ is non-empty by observing that $\mathbf{F}^* \in \mathcal{R}_X(\mathbf{d})$ provided that $p_{\mathbf{F}^0}$ does not ramify everywhere and such that $p_{\mathbf{F}^j}$ does not ramify everywhere along any non-empty irreducible component of $\text{ram}(p_{\mathbf{F}^{j-1}}) \cap \dots \cap \text{ram}(p_{\mathbf{F}^0}) \cap U$, for each j with $0 < j \leq m$.

The asserted $\mathfrak{B}(\mathbf{d})_e$ is the intersection with $\mathfrak{R}_X(\mathbf{d})$ of the Zariski closed subset

$$\mathfrak{B}'(\mathbf{d})_e \subset \mathcal{U}_X(\mathbf{d})$$

constructed as the union over j , $0 \leq j \leq m$, of closed subsets

$$\mathcal{U}_X(d_0) \times \dots \times \mathcal{U}_X(d_{j-1}) \times \mathcal{B}(d_0, \dots, d_j)_e \times \dots \times \mathcal{U}_X(d_m)$$

for suitably defined closed subsets $\mathcal{B}(d_0, \dots, d_j)_e \subset \mathcal{U}_X(d_j)$. Namely, $\mathcal{B}(d_0)_e \subset \mathcal{U}_X(d_0)$ is the closed subset given by Proposition 1.3 for d_0, e . By Lemma 1.6, the residual cycles $R_{\mathbf{F}}(Z)$ constructed for $\mathbf{F}_0 \in \mathcal{U}_X(d_0)$ and effective cycles Z on X of degree bounded by e are of degrees bounded by some e_1 (depending upon both e and d_0). We define

$\mathcal{B}(d_0, d_1)_e \subset \mathcal{U}_X(d_1)$ to be the closed subset given by Proposition 1.3 for d_1, e_1 . Continuing inductively, we conclude by Lemma 1.6 that the residual cycles $R_{\mathbf{F}_{j-1}} \circ R_{\mathbf{F}_0}(Z)$ constructed for $\mathbf{F}_0 \in \mathcal{U}_X(d_0), \dots, \mathbf{F}_{j-1} \in \mathcal{U}_X(d_{j-1})$ and effective cycles Z on X of degree bounded by e are of degrees bounded by some e_j (depending upon e and d_0, \dots, d_{j-1}). We define $\mathcal{B}(d_0, \dots, d_j)_e \subset \mathcal{U}_X(d_j)$ to be the closed subset given by Proposition 1.3 for d_j, e_j .

By Corollary 1.5, any component of excess dimension of $|Y| \cap |R_{\mathbf{F}^*}(Z)|$ must be contained in $|Y| \cap |Z| \cap \text{ram}(p_{\mathbf{F}^m}) \cap \dots \cap \text{ram}(p_{\mathbf{F}^0})$ provided that $\mathbf{F}^* \notin \mathcal{B}'(\mathbf{d})_e$. Thus, for $\mathbf{F}^* \in \mathfrak{R}_X(\mathbf{d}), \mathbf{F}^* \notin \mathcal{B}'(\mathbf{d})_e$, any component of excess dimension of $|Y| \cap |R_{\mathbf{F}^*}(Z)|$ must be contained in $\text{ram}(p_{\mathbf{F}^m}) \cap \dots \cap \text{ram}(p_{\mathbf{F}^0}) \subset X - U$.

By Proposition 1.3, for any integer c we may choose each d_j (depending as above on e and the d_i 's for $i < j$) so that $\mathcal{B}(d_0, \dots, d_j)_e \subset \mathcal{U}_X(d_j)$ has codimension at least c . This insures that

$$\mathfrak{B}(\mathbf{d})_e \equiv \mathfrak{B}'(\mathbf{d})_e \cap \mathfrak{R}(\mathbf{d}) \subset \mathfrak{R}(\mathbf{d})$$

also has codimension at least c . \square

Note 1.8. Note from the paragraph above that each d_j may be chosen *arbitrarily*, so long as it is sufficiently large. This lower bound depends only on c, e and the choice of the previous integers d_1, \dots, d_{j-1} .

In §3 we will use Theorem 1.7 in conjunction with the following proposition which expresses a cycle Z in terms of projecting cones and the cycle $R_{\mathbf{F}^*}(Z)$. This proposition follows immediately from Corollary 1.5 and Lemma 1.6.

Proposition 1.9. *Let $X \subset \mathbf{P}^n$ be a closed subvariety of dimension m , let r, s be non-negative integers with $r + s \geq m$, and let e be a positive integer. Assume that \mathbf{d} is chosen so that the codimension of $\mathcal{B}(\mathbf{d})_e \subset \mathcal{R}_X(\mathbf{d})$ is at least 1 and that the field k is sufficiently large that there exists a k -rational point $\mathbf{F}^* \in \mathfrak{R}_X(\mathbf{d}) - \mathfrak{B}(\mathbf{d})_e$. Choose some such \mathbf{F}^* . Consider the equality*

$$Z = \pi_{\mathbf{F}^0}^* \{p_{\mathbf{F}^0}^*(Z)\} \bullet X + \sum_{i=1}^m (-1)^i \pi_{\mathbf{F}^i}^* \{p_{\mathbf{F}^i}^* \{R_{\mathbf{F}^{i-1}} \circ \dots \circ R_{\mathbf{F}^0}(Z)\}\} \bullet X + (-1)^{m+1} R_{\mathbf{F}^*}(Z) \quad (1.9.1)$$

and let $\psi_{\mathbf{F}^*}^+(Z), \psi_{\mathbf{F}^*}^-(Z)$ denote the positive and negative parts of the right hand side of (1.9.1). Then:

- (a) For each effective s -cycle Z on X , $(\psi_{\mathbf{F}^*}^+(Z), \psi_{\mathbf{F}^*}^-(Z))$ is a pair of effective s -cycles on X with the property that

$$Z = \psi_{\mathbf{F}^*}^+(Z) - \psi_{\mathbf{F}^*}^-(Z).$$

- (b) For all effective cycles Y, Z on X of dimension r, s which are of degrees $\leq e$, any component of excess dimension of either $|Y| \cap |\psi_{\mathbf{F}^*}^+(Z)|$ or $|Y| \cap |\psi_{\mathbf{F}^*}^-(Z)|$ lies in $|Y| \cap |Z|$.

- (c) This construction determines a continuous algebraic map

$$\psi_{\mathbf{F}^*} = (\psi_{\mathbf{F}^*}^+, \psi_{\mathbf{F}^*}^-) : \mathcal{C}_s(X) \rightarrow \mathcal{C}_s(X)^2.$$

Moreover, there is an integer e' such that

$$\psi_{\mathbf{F}^*}(\mathcal{C}_{s, \leq e}(X)) \subset \mathcal{C}_{s, \leq e'}(X)^2.$$

2. Moves in Projective Space

In this section, we consider effective r -cycles Y and effective s -cycles Z on \mathbf{P}^m of degree bounded by a fixed positive integer e , where $r + s \geq m$. We present a construction on all s -cycles on \mathbf{P}^m which moves each such Z so that the resulting cycle intersects properly each such Y . The method we employ generalizes a technique introduced in [L] to move s -cycles Z of bounded degree to intersect properly a given hyperplane $L \equiv \mathbf{P}^{m-1} \subset \mathbf{P}^m$. One can interpret this method as taking the “algebraic suspension” $z_1 \# Z \subset \mathbf{P}^{m+1} = z_0 \# \mathbf{P}^m$ and observing that $z_1 \# Z$ intersects $z_0 \# L$ properly provided that $z_1 \neq z_0$. One then observes that for most divisors $D \subset \mathbf{P}^{m+1}$ of sufficiently high degree the projection of $(z_1 \# Z) \bullet D$ off z_0 meets L properly.

Throughout this section, we consider non-negative integers $r, s \leq m$ with $r + s \geq m$. We set $t \equiv m - r$. We shall fix a linear embedding

$$\mathbf{P}^m \subset \mathbf{P}^{m+t}, \quad (m + t = 2m - r)$$

and some fixed linear subspace $L_0 \subset \mathbf{P}^{m+t}$ of dimension $t - 1$ missing \mathbf{P}^m . Thus,

$$\mathbf{P}^{m+t} = L_0 \# \mathbf{P}^m,$$

where $\#$ denotes the “algebraic join”.

We next introduce our “parameter space of moves” for s -cycles in \mathbf{P}^m .

Definition 2.1. For any t -tuple of positive integers $\mathbf{N} = (N_1, \dots, N_t)$, let

$$\mathcal{L}(\mathbf{N}) \subset \text{Grass}_{t-1}(\mathbf{P}^{m+t}) \times \prod_{i=1}^t \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^{m+t}}(N_i)))$$

denote the open set consisting of those (L, \mathbf{D}) , where $\mathbf{D} = (D_1, \dots, D_t)$ which satisfy

- (a) $L \cap \mathbf{P}^m = \emptyset = L \cap L_0$,
- (b) $|D_1| \cap \dots \cap |D_t| \cap L = \emptyset = |D_1| \cap \dots \cap |D_t| \cap L_0$.

For any $L \in \text{Grass}_{t-1}(\mathbf{P}^{m+t})$, we denote by $\mathcal{L}_L(\mathbf{N})$ the Zariski dense open subset of $\prod_{i=1}^t \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^{m+t}}(N_i)))$ given as the fibre of the projection $\mathcal{L}(\mathbf{N}) \rightarrow \text{Grass}_{t-1}(\mathbf{P}^{m+t})$ above L .

Observe that for any k -rational point $(L, \mathbf{D}) \in \mathcal{L}(\mathbf{N})$, the intersection product $D_1 \bullet \dots \bullet D_t$ is a well defined n -cycle on \mathbf{P}^{m+t} with support $|D_1| \cap \dots \cap |D_t|$ which meets $L \# Z$ properly and which misses L_0 , the center of the projection π_0 .

In the following proposition, we establish a constructible set of arbitrarily high codimension of “good moves” for effective s -cycles Z of degree $\leq e$ which satisfy a single intersection condition depending upon a chosen $(t - 1)$ - dimensional linear subspace L . We recall that a subset of a quasi-projective variety is said to be **constructible** if it can be written as a finite disjoint union of (Zariski) locally-closed subsets. The key property of constructible subsets is that any map $f : X \rightarrow Y$ of quasi-projective varieties sends a constructible subset $C \subset X$ to a constructible subset $f(C) \subset Y$.

Proposition 2.2. *Let L be a $(t - 1)$ -dimensional linear subspace of \mathbf{P}^{m+t} satisfying $L \cap \mathbf{P}^m = \emptyset = L \cap L_0$. For a given positive integer e and a t -tuple of positive integers $\mathbf{N} = (N_1, \dots, N_t)$, there exists a constructible subset*

$$\mathcal{B}_L(\mathbf{N})_e \subset \mathcal{L}_L(\mathbf{N})$$

such that

- (a) For a constant K depending only upon X and e ,

$$\text{codim} \mathcal{B}_L(\mathbf{N})_e \geq \min(N_i) - K,$$

- (b) For every k -rational point $\mathbf{D} = (D_1, \dots, D_t) \notin \mathcal{B}_L(\mathbf{N})_e$, the cycles

$$|L_0 \# Y| \cap |L \# Z| \cap |D_1| \cap \dots \cap |D_{i-1}| \quad \text{and} \quad |D_i|$$

meet properly for all $i \geq 0$ whenever Y is an effective r -cycle and Z is an effective s -cycle both of degree $\leq e$ on \mathbf{P}^m with the property that $L_0 \# Y$ intersects $L \# Z$ properly.

Proof. Let

$$\mathcal{J} \subset \mathcal{C}_{r, \leq e}(\mathbf{P}^m) \times \mathcal{C}_{s, \leq e}(\mathbf{P}^m) \times \mathbf{P}^{m+t}$$

denote the incidence correspondence of triples (Y, Z, x) satisfying $x \in |L_0 \# Y| \cap |L \# Z|$. We define

$$\mathcal{G}_{L,e} \subset \mathcal{C}_{r, \leq e}(\mathbf{P}^m) \times \mathcal{C}_{s, \leq e}(\mathbf{P}^m)$$

to be the open subset above which the fibre of the projection

$$\mathcal{J} \rightarrow \mathcal{C}_{r, \leq e}(\mathbf{P}^m) \times \mathcal{C}_{s, \leq e}(\mathbf{P}^m)$$

has proper dimension (i.e., $r + s - m + t = s$). Note that $\mathcal{G}_{L,e}$ consists of pairs of cycles (Y, Z) such that $L_0 \# Y$ and $L \# Z$ intersect properly. We consider the closed subset

$$\tilde{\mathcal{B}}_L(\mathbf{N})_e \subset \mathcal{G}_{L,e} \times \mathcal{L}_L(\mathbf{N})$$

consisting of those tuples (Y, Z, \mathbf{D}) satisfying the condition that $L_0 \# Y$ and $L \# Z$ intersect properly but

$$|L_0 \# Y| \cap |L \# Z| \cap |D_1| \cap \dots \cap |D_{i-1}| \quad , \quad |D_i|$$

intersect improperly for some $i \leq t$. Finally, we set

$$\mathcal{B}_L(\mathbf{N})_e = \text{pr}_{3*}[(\tilde{\mathcal{B}}_L)(\mathbf{N})_e].$$

We proceed to verify that the fibre $\tilde{\mathcal{B}}_{Y,Z} \subset \mathcal{L}_L(\mathbf{N})$ of $\tilde{\mathcal{B}}_L(\mathbf{N})_e$ above any $(Y, Z \in \mathcal{G}_{L,e})$ has codimension at least N if each $N_i \geq N$. This will immediately imply the codimension of $\mathcal{B}_L(\mathbf{N})_e \subset \mathcal{L}_L(\mathbf{N})$ is greater than some prescribed positive integer c provided that

$$N \geq c + \dim(\mathcal{C}_{r,\leq e}(\mathbf{P}^m)) + \dim(\mathcal{C}_{s,\leq e}(\mathbf{P}^m)).$$

The condition that $W_j \equiv |L_0 \# Y| \cap |L \# Z| \cap |D_1| \cap \dots \cap |D_{j-1}|$ meet $|D_j|$ improperly is equivalent to the condition that the restriction map

$$\mathcal{O}_{\mathbf{P}^{m+t}}(N_j) \longrightarrow \mathcal{O}_{W_j}(N_j) \longrightarrow \mathcal{O}_{W_{j,\omega}}(N_j)$$

sends $D_j \in \Gamma(\mathcal{O}_{\mathbf{P}^{m+t}}(N_j))$ to 0 for some irreducible component $W_{j,\omega} \subset W_j$. Since $W_{j,\omega}$ has positive dimension, this composition has rank at least N . To see this note that if we restrict to some affine open $\mathbf{A}^m \subset \mathbf{P}^m$ containing a dense open subset of $W_{j,\omega}$, then the dimension of the subspace of regular functions on $\mathbf{A}^m \cap W_{j,\omega}$ of degree $\leq N$ is at least N . We conclude that the closed subvariety $\tilde{\mathcal{B}}_{Y,Z,j} \subset \mathcal{L}_L(\mathbf{N})$ consisting of those \mathbf{D} with D_j meeting W_j improperly and $D_{j'}$ meeting $W_{j'}$ properly for $j' < j$ has codimension at least N . Thus,

$$\tilde{\mathcal{B}}_{Y,Z} = \coprod_j \tilde{\mathcal{B}}_{Y,Z,j} \subset \mathcal{L}_L(\mathbf{N})$$

also has codimension at least N . \square

Proposition 2.2 suggests the following construction of moving cycles in \mathbf{P}^m , sending the s -cycle Z to $\varphi_{L,\mathbf{D}}(Z)$.

Proposition 2.3 *Let $\pi_0 : \mathbf{P}^{m+t} \dashrightarrow \mathbf{P}^m$ denote the linear projection with center L_0 . For any $e > 0$ and any t -tuple $\mathbf{N} = (N_1, \dots, N_t)$, the assignment*

$$\varphi_{L,\mathbf{D}}(Z) \stackrel{\text{def}}{=} \pi_{0*}((L \# Z) \bullet D_1 \bullet \dots \bullet D_t)$$

determines a continuous algebraic map

$$\Phi : \mathcal{C}_s(\mathbf{P}^m) \times \mathcal{L}(\mathbf{N}) \longrightarrow \mathcal{C}_s(\mathbf{P}^m)$$

with $\Phi(\mathcal{C}_{s,\leq e}(\mathbf{P}^m) \times \mathcal{L}(\mathbf{N})) \subset \mathcal{C}_{s,\leq e|\mathbf{N}|}(\mathbf{P}^m)$, where $|\mathbf{N}| \stackrel{\text{def}}{=} \prod_i N_i$. This construction has the following properties:

- (a) For any k -rational point $(L, \mathbf{D}) \in \mathcal{L}(\mathbf{N})$ and any effective s -cycle Z on \mathbf{P}^m , $\varphi_{L,\mathbf{D}}(Z)$ is also an effective s -cycle on \mathbf{P}^m .
- (b) For any k -rational point $(L, \mathbf{D}) \in \mathcal{L}(\mathbf{N})$ with the property that $D_1 \bullet \dots \bullet D_t = |\mathbf{N}| \cdot \mathbf{P}^m$,

$$\varphi_{L,\mathbf{D}}(Z) = |\mathbf{N}| \cdot Z$$

for all effective s -cycles Z .

- (c) If (L, \mathbf{D}) is a k -rational point of $\mathcal{L}(\mathbf{N})$ such that $\mathbf{D} \notin \mathcal{B}_L(\mathbf{N})_e$ (cf. (2.2)), if Y, Z are effective cycles on \mathbf{P}^m of dimension r, s and degree $\leq e$, and if $L_0 \# Y$ meets $L \# Z$ properly, then

$$Y \quad , \quad \varphi_{L, \mathbf{D}}(Z)$$

intersect properly.

Proof. If Z on \mathbf{P}^m is an s -effective cycle of degree d , then $L \# Z$ is an effective $(s+t)$ -cycle \mathbf{P}^{m+t} also of degree d . Consequently, $(L \# Z) \bullet D_1 \bullet \dots \bullet D_t$ and thus also $\varphi_{L, \mathbf{D}}(Z)$ have degree $d \cdot |\mathbf{N}|$. Moreover, sending $\langle Z, (L, \mathbf{D}) \rangle$ to $\varphi_{L, \mathbf{D}}(Z)$ determines a continuous algebraic map, for Φ is a composition of operations each of which is given by a continuous algebraic map (cf. [F]).

As observed following Definition 2.1, the intersection product $D_1 \bullet \dots \bullet D_t$ is a well defined n -cycle on \mathbf{P}^{m+t} with support $|D_1| \cap \dots \cap |D_t|$ which meets $L \# Z$ properly and which misses L_0 for any k -rational point $(L, \mathbf{D}) \in \mathcal{L}(\mathbf{N})$. Thus, $\varphi_{L, \mathbf{D}}(Z)$ is a well defined s -cycle on \mathbf{P}^m .

To prove that $\varphi_{L, \mathbf{D}}(Z) = |\mathbf{N}| \cdot Z$ whenever $D_1 \bullet \dots \bullet D_t = |\mathbf{N}| \cdot \mathbf{P}^m$, observe that $\mathbf{P}^m \subset \mathbf{P}^{m+t}$ meets $L \# Z$ properly (since $L \cap \mathbf{P}^m = \emptyset$) and transversely at all regular points of Z .

If $(L, \mathbf{D}) \in \mathcal{L}(\mathbf{N})$, then $L_0 \# Y, L \# Z$ have pure dimension $m, s+t$ respectively. The hypothesis that $L_0 \# Y$ meets $L \# Z$ properly thus implies that $|L_0 \# Y| \cap |L \# Z|$ has dimension s . We thus may apply Proposition 2.2 to conclude that if $\mathbf{D} \notin \mathcal{B}(\mathbf{N})_e$, if Y, Z are effective cycles of degrees $\leq e$, and if $L_0 \# Y$ meets $L \# Z$ properly, then $(L_0 \# Y) \bullet (L \# Z) \bullet D_1 \bullet \dots \bullet D_t$ has pure dimension $r + s - m$. Because $L_0 \cap |D_1| \cap \dots \cap |D_t| = \emptyset$, π_{0*} restricts to a (well defined) finite map on $|(L \# Z) \bullet D_1 \bullet \dots \bullet D_t|$. We apply the projection formula to π_{0*} to conclude that

$$Y \bullet \pi_{0*}[(L \# Z) \bullet D_1 \bullet \dots \bullet D_t] = \pi_{0*}[(L_0 \# Y) \bullet (L \# Z) \bullet D_1 \bullet \dots \bullet D_t].$$

Hence, $Y, \varphi_{L, \mathbf{D}}(Z)$ meet properly. \square

The following lemma will enable us to apply Proposition 2.3 to all effective cycles of bounded degree.

Lemma 2.4. *Let L be a k -rational point of $\text{Grass}_{t-1}(\mathbf{P}^{m+t})$ satisfying $L \cap P^m = \emptyset = L \cap L_0$. Then*

$$(L_0 \# Y), (L \# Z) \text{ intersect properly}$$

in \mathbf{P}^{m+t} for every r -cycle Y, s -cycle Z on \mathbf{P}^m such that either $|L_0 \# Y| \cap L = \emptyset$ or Y intersects Z properly.

Proof. We first check that if Y, Z intersect properly, so do $(L_0 \# Y), (L \# Z)$ for any choice of L . Observe that

$$|L_0 \# Y| \cap P^m = |Y|, \quad |L \# Z| \cap P^m = |Z|.$$

Thus, if some component of $|L_0 \# Y| \cap |L \# Z|$ has dimension $> s$, then some component of $|L_0 \# Y| \cap |L \# Z| \cap P^m = |Y| \cap |Z|$ would have dimension $> s - t$.

On the other hand, if $|L_0 \# Y| \cap L = \emptyset$, then $L \# z$ intersects $L_0 \# Y$ in only finitely many points for any $z \in \mathbf{P}^m$. Thus, $|L_0 \# Y| \cap |L \# Z|$ has dimension at most s . \square

Combining Proposition 2.3 and Lemma 2.4, we now verify for any given e that we can “move” all effective s -cycles Z of degree $\leq e$ so that the resulting cycles intersect properly every effective r -cycle Y of degree $\leq e$.

As in Proposition 1.9, we require the existence of k -rational points avoiding certain “bad” closed subsets of positive codimension. The existence of such k -rational points is automatic for an infinite field, but for a finite field all rational points of a given variety might lie on a subvariety of positive codimension. On the other hand, if k is a finite field and if there are only finitely many “bad” subvarieties of positive codimension, there will exist some finite extension k' of k such that there exist k' -rational points avoiding these “bad sets”. Moreover, any extension k'' of k' will evidently have this same property.

Since our definitions of “good” and “bad” sets are independent of k , we conclude that all “sufficiently large” finite fields k of a given residue characteristic admit the existence of k -rational points missing the “bad” sets. We shall use the expression “ k is a **sufficiently large finite field**” to mean that there exist k -rational points avoiding the bad sets (of positive codimension) occurring in our constructions.

Proposition 2.5. *Fix positive integers e and c . Assume that either k is an infinite field or a sufficiently large finite field. Consider a sequence $\mathbf{N}_1, \mathbf{N}_2, \mathbf{N}_3, \dots$ of t -tuples of positive integers, and set $e_1 = e$ and $e_{j+1} = |\mathbf{N}_j|e_j$ for $j > 1$. Assume that the \mathbf{N}_j 's are chosen so that $\min_{1 \leq i \leq t} N_{j+1,i} > K(e_j) + c$ where $K(e_j)$ is the constant in Proposition 2.2a. Then there exist an integer $E \leq \dim \mathcal{C}_{s, \leq e}(\mathbf{P}^m)$ and $(t-1)$ -dimensional linear subspaces L_1, \dots, L_E of \mathbf{P}^{m+t} , each missing L_0 and \mathbf{P}^m , with the following properties:*

(a) For each j ,

$$\text{codim} \left\{ \mathcal{L}_{L_j}(\mathbf{N}_j) - \mathcal{B}_{L_j}(\mathbf{N}_j)_{e_j} \right\} \geq c,$$

(b) For any k -rational point

$$(\mathbf{D}_1, \dots, \mathbf{D}_E) \in \prod_{j=1}^E \mathcal{L}_{L_j}(\mathbf{N}_j) - \mathcal{B}_{L_j}(\mathbf{N}_j)_{e_j}$$

and any effective cycles Y, Z on \mathbf{P}^m of dimensions r, s and degrees $\leq e$,

$$Y \quad , \quad \varphi_{L_E, \mathbf{D}_E} \circ \dots \circ \varphi_{L_1, \mathbf{D}_1}(Z)$$

intersect properly.

Proof. Note to begin that assertion (a) follows immediately from Proposition 2.2a, so it remains to prove assertion (b).

For each (of the finitely many) irreducible components A_γ of $A = C_{r, \leq e}(\mathbf{P}^m)$ of maximal dimension d_e , choose some k -rational point $a(\gamma) \in A_\gamma$ (possible since k is assumed either infinite or sufficiently large). Using once again the fact that k is infinite or sufficiently

large, we may choose some $(t - 1)$ -dimensional linear plane L_1 in \mathbf{P}^{m+t} missing P^m, L_0 , and each $L_0 \# Y_{a(\gamma)}$. Let $A_1 \subset A$ denote the closed subvariety of those Y 's such that $L_1 \cap |L_0 \# Y| \neq \emptyset$, and observe that each component of A_1 has dimension strictly less than d_e . By Proposition 2.3 and Lemma 2.4, Y and $\varphi_{L_1, \mathbf{D}_1}(Z)$ intersect properly for any effective r -cycle Y of degree $\leq e$ and any effective s -cycle Z of degree $\leq e$ which does not lie in A_1 whenever \mathbf{D}_1 is k -rational point of $\mathcal{L}_{L_1}(\mathbf{N}_1) - \mathcal{B}_{L_1}(\mathbf{N}_1)_{e_1}$.

Proceeding inductively, assume for some $j \geq 1$ that we have defined the closed subvariety $A_j \subset A$ of codimension at least j which satisfies the condition that

$$Y, \varphi_{L_j, \mathbf{D}_j} \circ \cdots \circ \varphi_{L_1, \mathbf{D}_1}(Z)$$

intersect properly for any effective r -cycle Y of degree $\leq e$ and any effective s -cycle Z of degree $\leq e$ which does not lie in A_j . For each (of the finitely many) irreducible components $A_{j, \gamma}$ of maximal dimension choose as above some point $a(\gamma) \in A_{j, \gamma}$ and choose L_{j+1} missing P^m, L_0 , and each $L_0 \# Y_{a(\gamma)}$. We set $A_{j+1} \subset A_j$ equal to the closed subvariety of that Y 's such that $L_{j+1} \cap |L_0 \# Y| = \emptyset$. Applying Proposition 2.3 and Lemma 2.4 once again, we conclude that

$$Y, \varphi_{L_j, \mathbf{D}_j} \circ \cdots \circ \varphi_{L_1, \mathbf{D}_1}(Z)$$

intersect properly for any effective r -cycle Y of degree $\leq e$ and any effective s -cycle Z of degree $\leq e$ which does not lie in A_j , whenever \mathbf{D}_j is k -rational point of $\mathcal{L}_{L_j}(\mathbf{N}_j) - \mathcal{B}_{L_j}(\mathbf{N}_j)_{e_j}$.

Finally, we see that this process stops after E steps, where $E \leq d_e + 1$. \square

The following theorem is our ‘‘Moving Lemma for Cycles of Bounded Degree’’ in the special case in which the variety X is projective space \mathbf{P}^m itself.

Theorem 2.6. *Fix positive integers e and c . Assume that either k is an infinite field or a sufficiently large finite field. Choose $\mathbf{N}_1, \dots, \mathbf{N}_E$ and L_1, \dots, L_E as in Proposition 2.5 and let*

$$\underline{0} \stackrel{\text{def}}{=} (\mathbf{D}_1, \dots, \mathbf{D}_E) \in \prod_{j=1}^E \mathcal{L}_{L_j}(\mathbf{N}_j)$$

be a k -rational point such that $D_{j,1} \bullet \cdots \bullet D_{j,t} = |\mathbf{N}_j| \cdot \mathbf{P}^m$ for each $j, 1 \leq j \leq E$. (For example, suppose that $D_{j,i} = N_{j,i} H_{j,i}$ where $H_{j,1}, \dots, H_{j,t}$ are k -rational hyperplanes in general position.) Let

$$\ell \subset \prod_{j=1}^E \prod_{i=1}^t \mathbf{P}(\Gamma(\mathcal{O}_{\mathbf{P}^{m+t}}(N_{j,i})))$$

be any k -rational line through $\underline{0}$ chosen so that

$$\ell^\circ \equiv \ell \cap \prod_{j=1}^E \mathcal{L}_{L_j}(\mathbf{N}_j) - \mathcal{B}_{L_j}(\mathbf{N}_j)_{e_j} \subset \ell$$

is dense. Then there exists a continuous algebraic map

$$\Theta : \mathcal{C}_s(\mathbf{P}^m) \times \ell^\circ \rightarrow \mathcal{C}_s(\mathbf{P}^m)$$

with the following properties. Set $\theta_p = \Theta|_{\mathcal{C}_s(\mathbf{P}^m) \times \{p\}}$.

- (a) $\theta_{\underline{0}}(Z) = \prod_{j=1}^E |\mathbf{N}_j| \cdot Z$ for any effective s -cycle Z .
- (b) For any k -rational point $p \in \ell^\circ$, θ_p determines a linear map on effective s -cycles on \mathbf{P}^m .
- (c) For any effective s -cycle Z , the restriction of Θ to $\{Z\} \times \ell^\circ$ determines a rational equivalence $Z_\theta \subset \mathbf{P}^m \times \ell$ whose fibre above a point $p \in \ell^\circ$ is the cycle with Chow point $\theta_p(Z)$.
- (d) For all effective cycles Y, Z on \mathbf{P}^m of dimensions r, s and degrees $\leq e$ and all k -rational points $p \in \ell^\circ - \underline{0}$,

$$Y, \theta_p(Z)$$

intersect properly.

Proof. Iterating the continuous algebraic map Φ of Proposition 2.3, we obtain the continuous algebraic map

$$\tilde{\Theta} : \mathcal{C}_s(\mathbf{P}^m) \times \mathcal{L}(\mathbf{N}_1) \times \cdots \times \mathcal{L}(\mathbf{N}_1) \rightarrow \mathcal{C}_s(\mathbf{P}^m).$$

We define Θ to be the restriction of $\tilde{\Theta}$ to $\mathcal{C}_s(\mathbf{P}^m) \times \ell^\circ$.

Property (a) is given by Proposition 2.3.b, whereas property (b) follows from Proposition 2.3.a. To verify property (c), we let λ denote the generic point of ℓ and $k(\lambda)$ denote the function field of ℓ . Then $\Theta(Z, \lambda)$ determines a cycle on \mathbf{P}^m rational over some radicial extension of $k(\lambda)$ (cf. [F]). On the other hand, since the operations used to define Θ send cycles to cycles, we see that the cycle Z_λ with Chow point $\Theta(Z, \lambda)$ is in fact a cycle on $\mathbf{P}_{k(\lambda)}^m$. The closure of this cycle, Z_θ on $\mathbf{P}^m \times \ell$, is the asserted rational equivalence. (Observe that Z_θ is necessarily flat over ℓ so that the fibre $p \in \ell$ is given by specialization of Z_λ .) Finally, property (d) follows from Proposition 2.5. \square

3. Main Theorem and Consequences

Using Theorem 1.7, Proposition 1.9, and Theorem 2.6, we now prove our “Moving Lemma for Cycles of Bounded Degree”.

Theorem 3.1. *Let $X \subset \mathbf{P}^n$ be a projective variety of dimension m . Assume that either k is an infinite field or a sufficiently large finite field. Let r, s, e be non-negative integers with $r + s \geq m$. Then there exists a Zariski open neighborhood $\mathcal{O} \subset \mathbf{P}^1$ of a k -rational point $0 \in \mathbf{P}^1$ and a continuous algebraic map*

$$\Psi = (\Psi^+, \Psi^-) : \mathcal{C}_s(X) \times \mathcal{O} \rightarrow \mathcal{C}_s(X)^2$$

satisfying the following properties (where $\psi_p^\pm = \Psi^\pm|_{\mathcal{C}_s(X) \times \{p\}}$):

- (a) For some positive integer M and some k -rational point $\mathbf{F}^* \in \mathfrak{R}_X(d) - \mathfrak{B}(d)_e$, chosen as in 1.9, one has

$$\psi_0^+ = (M+1) \cdot \psi_{\mathbf{F}^*}^+ + M \cdot \psi_{\mathbf{F}^*}^- \quad , \quad \psi_0^- = (M+1) \cdot \psi_{\mathbf{F}^*}^- + M \cdot \psi_{\mathbf{F}^*}^+$$

where $\psi_{\mathbf{F}^*}^+, \psi_{\mathbf{F}^*}^- : \mathcal{C}_s(X) \rightarrow \mathcal{C}_s(X)$ are the continuous algebraic maps of Proposition 1.9.a. In particular, for every effective s -cycle Z ,

$$Z = \psi_0^+(Z) - \psi_0^-(Z).$$

- (b) For any k -rational point $p \in \mathcal{O}$, ψ_p^\pm determines a pair of linear maps on effective s -cycles
- (c) For any effective s -cycle Z on X , the restriction of Ψ to $\{Z\} \times \mathcal{O}$ determines a rational equivalence $Z_\psi \subset \mathbf{P}^n \times \mathbf{P}^1$ whose fibre above a point $p \in \mathcal{O}$ is $\psi_p^+(Z) - \psi_p^-(Z)$.
- (d) For all effective cycles Y, Z on X of dimensions r, s and degrees $\leq e$ and all k -rational points $p \in \mathcal{O} - \{0\}$, any component of excess dimension (i.e., $> r+s-m$) of either $|Y| \cap |\psi_p^+(Z)|$ or $|Y| \cap |\psi_p^-(Z)|$ lies in the singular locus of X .

Proof. Choose \mathbf{F}^* as in Proposition 1.9 and recall the equality (1.9.1)

$$Z = \pi_{\mathbf{F}^*}^* \{p_{\mathbf{F}^*}^0(Z)\} \bullet X + \sum_{i=1}^m (-1)^i \pi_{\mathbf{F}^*}^* \{p_{\mathbf{F}^*}^i \{R_{\mathbf{F}^*}^{i-1} \circ \cdots \circ R_{\mathbf{F}^*}^0(Z)\}\} \bullet X + (-1)^{m+1} R_{\mathbf{F}^*}^m(Z)$$

and the fact that $\psi_{\mathbf{F}^*}^\pm$ are given as the positive and negative parts of the right hand side of (1.9.1). Let $e' \geq e$ be a positive integer bounding the degrees of $p_{\mathbf{F}^*}^i(Y)$ and of $p_{\mathbf{F}^*}^i(R_{\mathbf{F}^*}^{i-1} \circ \cdots \circ R_{\mathbf{F}^*}^0(Z))$ as Y and Z vary over effective r -cycles and s -cycles on X (respectively) of degrees $\leq e$, and i varies from 1 to m .

We now apply Theorem 2.6 with e replaced by e' and with some fixed $c > 1$. In fact we apply the theorem twice. We choose two sequences $\mathbf{N} = (\mathbf{N}_1, \dots, \mathbf{N}_E)$ and $\mathbf{N}' = (\mathbf{N}'_1, \dots, \mathbf{N}'_{E'})$ with $N_{j,i}$ and $N'_{j,i}$ sufficiently large as required in Proposition 2.5, so that

$$\prod_{j=1}^E |\mathbf{N}_j| = M+1 \quad , \quad \prod_{j=1}^{E'} |\mathbf{N}'_j| = M$$

for some positive integer M . To see that this is possible recall that the integers $N_{j,i}$ can be chosen arbitrarily subject to the inequalities of 2.5. Consequently we may choose the $N_{j,i}$'s and the $N'_{j,i}$ so that $\prod_j |\mathbf{N}_j|$ and $\prod_j |\mathbf{N}'_j|$ are relatively prime. Then there exist positive integers a and a' so that $a \prod_j |\mathbf{N}_j| - a' \prod_j |\mathbf{N}'_j| = \pm 1$. We may assume, by reordering the choice if necessary, that the difference is $+1$, and we then replace $N_{E,t}$ with $aN_{E,t}$ and $N'_{E',t}$ with $a'N'_{E',t}$.

By Proposition 2.5 we obtain sequences (L_1, L_2, \dots, L_E) and $(L'_1, L'_2, \dots, L'_{E'})$ of $(t-1)$ -dimensional linear subspaces, for which the conclusions of 2.5 hold.

We now apply Theorem 2.6 (twice) for the $\{\mathbf{N}_j\}, \{L_j\}$ (and the $\{\mathbf{N}'_j\}, \{L'_j\}$) chosen above. We fix k -rational points $\underline{0}$ and $\underline{0}'$ and k -rational lines ℓ and ℓ' as in Theorem 2.6, and we let

$$\Theta_{\underline{\mathbf{N}}} : \mathcal{C}_s(\mathbf{P}^m) \times \ell^\circ \rightarrow \mathcal{C}_s(\mathbf{P}^m) \quad \text{and} \quad \Theta'_{\underline{\mathbf{N}'}} : \mathcal{C}_s(\mathbf{P}^m) \times (\ell')^\circ \rightarrow \mathcal{C}_s(\mathbf{P}^m)$$

be the continuous algebraic maps constructed in 2.6 for this data.

We now choose a dense open subset $\mathcal{O} \subset \mathbf{P}^1$ and point $0 \in \mathcal{O}$ together with open immersions $f : \mathcal{O} \hookrightarrow \ell^\circ$ and $f' : \mathcal{O} \hookrightarrow (\ell')^\circ$ sending 0 to $\underline{0}$ and $\underline{0}'$ respectively. We let $\Theta_{\underline{\mathbf{N}}, p}$ denote the restriction of $\Theta_{\underline{\mathbf{N}}}$ to $\mathcal{C}_s(\mathbf{P}^m) \times \{f(p)\}$ for any k -rational point $p \in \mathcal{O}$. We define $\Psi_{\underline{\mathbf{N}}}$ by

$$\Psi_{\underline{\mathbf{N}}}(Z, p) = (-1)^{m+1}(M+1) \cdot R_{\mathbf{F}^*}(Z) + \sum_{i=1}^m (-1)^i \pi_{\mathbf{F}^i}^* \left\{ \Theta_{\underline{\mathbf{N}}, p} \left\{ p_{\mathbf{F}^i*} (R_{\mathbf{F}^{i-1}} \circ \cdots \circ R_{\mathbf{F}^0}(Z)) \right\} \right\} \bullet X,$$

and define $\Psi_{\underline{\mathbf{N}'}}$ similarly. Finally, we define Ψ^\pm to be the positive and negative parts of $\Psi_{\underline{\mathbf{N}}} - \Psi_{\underline{\mathbf{N}'}}$, so that

$$\Psi^+(Z, p) - \Psi^-(Z, p) = \Psi_{\underline{\mathbf{N}}}(Z, p) - \Psi_{\underline{\mathbf{N}'}}(Z, p).$$

The fact that Ψ^+, Ψ^- are continuous algebraic maps follows immediately from Proposition 1.9 and Theorem 2.6. Property (a) is part of our definition of Ψ^\pm . Property (b) follows immediately from Proposition 1.9 and Theorem 2.6.b. The proof of property (c) is merely a repetition of the proof of Theorem 2.6.c.

To verify property (d), observe that Theorem 1.7.b implies that any component of excess dimension of $|Y| \cap |R_{\mathbf{F}^*}(Z)|$ lies in the singular locus of X . On the other hand, since $p_{\mathbf{F}^i}$ is a finite map,

$$Y \quad , \quad \pi_{\mathbf{F}^i}^* \left\{ \Theta_{\underline{\mathbf{N}}, p} \left\{ p_{\mathbf{F}^i*} (R_{\mathbf{F}^{i-1}} \circ \cdots \circ R_{\mathbf{F}^0}(Z)) \right\} \right\} \bullet X$$

intersect properly for any $i, 1 \leq i \leq m$, if and only if

$$p_{\mathbf{F}^i*}(Y) \quad , \quad \Theta_{\underline{\mathbf{N}}, p} \left\{ p_{\mathbf{F}^i*} (R_{\mathbf{F}^{i-1}} \circ \cdots \circ R_{\mathbf{F}^0}(Z)) \right\}$$

intersect properly. These do intersect properly whenever Y, Z are effective cycles of degrees $\leq e$ and $p \in \mathcal{O} - \{0\}$ by Theorem 2.6.d. The analogous remarks apply with \mathbf{N} replaced by \mathbf{N}' . This completes the proof. \square

Remark 3.2. The proof of Theorem 3.1 in fact proves the more general result that one can “move” effective s -cycles Z of degrees $\leq e$ to meet properly (off the singular locus of X) all effective cycles Y of dimension $\geq m - s$ and degree $\leq e$. Namely, we verify that the arguments of section 1 apply when $\mathcal{C}_{r, \leq e}(X)$ is replaced by

$$\mathcal{C}_{\geq n-s, \leq e}(X) \quad \equiv \quad \coprod_{r \geq m-s, d \leq e} \mathcal{C}_{r, d}(X).$$

Similarly, the arguments of section 2 apply when we replace $\mathcal{C}_{r,\leq e}(\mathbf{P}^n)$ by $\mathcal{C}_{\geq s,\leq e}(\mathbf{P}^n)$ and $t = n - r$ by s .

When $k = \mathbf{C}$ our Main Theorem can be phrased differently and in a way that may appeal to complex geometers. In this case the p -cycles form a topological abelian group where the topology is compactly generated by the images of $\mathcal{C}_{p,\leq e}(X) \times \mathcal{C}_{p,\leq e}(X)$ under the natural projection

$$\pi : \mathcal{C}_p(X) \times \mathcal{C}_p(X) \longrightarrow \mathcal{Z}_p(X)$$

We set $\mathcal{Z}_{p,\leq e}(X) \stackrel{\text{def}}{=} \pi\{\mathcal{C}_{p,\leq e}(X) \times \mathcal{C}_{p,\leq e}(X)\}$

Corollary 3.3. *Let $X \subset P_{\mathbf{C}}^n$ be a complex projective variety of dimension m . Let r, s, e be non-negative integers with $r + s \geq m$. Then there exist a Zariski open neighborhood \mathcal{O} of $\{0\}$ in \mathbf{C} and a continuous algebraic map*

$$\tilde{\Psi} : \mathcal{C}_s(X) \times \mathcal{O} \longrightarrow \mathcal{C}_s(X)^2$$

such that $\pi \circ \tilde{\Psi}$ induces by linearity a continuous map

$$\Psi : \mathcal{Z}_s(X) \times \mathcal{O} \longrightarrow \mathcal{Z}_s(X)$$

satisfying the following properties. Set $\psi_p = \Psi|_{\mathcal{Z}_s(X) \times \{p\}}$ for $p \in \mathcal{O}$.

- (a) $\psi_0 = \text{Id}$.
- (b) For any $Z \in \mathcal{Z}_s(X)$ and any $p \neq 0$ in \mathcal{O} , the restriction

$$\Psi|_{\{Z\} \times \mathcal{O}} : \{Z\} \times \mathcal{O} \longrightarrow \mathcal{Z}_s(X),$$

determines a rational equivalence between Z and $\psi_p(Z)$.

- (c) For any $p \in \mathcal{O}$, ψ_p is a continuous group homomorphism.
- (d) For any $Z \in \mathcal{Z}_{s,\leq e}(X)$, $Y \in$ and any $p \neq 0$ in \mathcal{O} , each component of excess dimension (i.e., $> r + s - m$) of the intersection

$$|Y| \cap |\psi_p(Z)|$$

is contained in the singular locus of X .

Remark. We may paraphrase Corollary 3.3 in terms of families of cycles. Let us say that a collection $\{Y_\alpha; \alpha \in A\}$ of r -cycles on X is a **collection of cycles of bounded degree** if there exists some positive integer e and some function $g : A \rightarrow \mathcal{Z}_{r,\leq e}(X)$ such that $Y_\alpha = g(\alpha)$ for all $\alpha \in A$. Then Corollary 3.3 asserts that for a given collection $\{Y_\alpha; \alpha \in A\}$ of r -cycles of bounded degree and a given collection $\{Z_\beta; \beta \in B\}$ of s -cycles of bounded degree, Ψ provides a continuous move of all s -cycles on X with the following property: for any $\alpha \in A, \beta \in B, p \in \mathcal{O} - \{0\}$ each component of $|Y_\alpha| \cap |\psi_p(Z_\beta)|$ of excess intersection lies in the singular locus of X .

Moreover, Corollary 3.3 further asserts how such a move is algebraic, thereby moving one “algebraic collection” to another.

As demonstrated in [R], [S], any s -cycle Z can be moved to intersect properly any r -cycle Y on a smooth, m -dimensional variety X with $r + s \geq m$. This does not immediately imply that the intersection product is well defined, for one must verify that if Z', Z'' are both obtained from Z by moving Z and both intersect Y properly, then $Y \bullet Z', Y \bullet Z''$ are rationally equivalent. As we see in the proof of the following theorem, the fact that such an intersection product is well defined is a straight-forward consequence of Theorem 3.1

Theorem 3.4 *Let X be a smooth, quasi-projective variety of dimension m and let Y, Z be cycles on X of dimension r, s with $r + s \geq m$. If both Z' and Z'' are rationally equivalent to Z and intersect Y properly, then $Y \bullet Z', Y \bullet Z''$ are also rationally equivalent.*

Proof. It suffices to assume that there exists some cycle Z_T on $T \times X$ equidimensional over a rational curve T with fibres $Z_0 = Z'$ and $Z_1 = Z''$. We first consider the case in which k is either an infinite field or a “sufficiently large” finite field as in the statement of Theorem 3.1. Choose some projective closure \bar{X} of X .

Write $Z_T = Z_T^+ - Z_T^-$, where Z_T^+, Z_T^- are flat families of effective cycles over T . (In other words, each of Z_T^+, Z_T^- are sums of irreducible subvarieties of $X \times T$ flat over T .) We may not assume that the fibres above 0, 1 of these flat families intersect Y properly, but we do know that those components of these fibres which do not intersect properly occur with the same multiplicity in positive and negative parts. The closures \bar{Z}_T^+, \bar{Z}_T^- of Z_T^+, Z_T^- are also flat families of effective cycles over T , thereby determining continuous algebraic maps (indeed, morphisms)

$$\varphi = (\varphi^+, \varphi^-) : T \rightarrow \mathcal{C}_s(\bar{X}) \times \mathcal{C}_s(\bar{X}).$$

We now apply Theorem 3.1 with e greater than the degrees of the closures of the positive and negative parts of Y as well as the degrees of \bar{Z}_t^+, \bar{Z}_t^- for all $t \in T$ to obtain Θ defined as the following composition

$$\sigma \circ (\tilde{\Psi} \times \tilde{\Psi}) \circ (\varphi \times \text{diag}) : T \times \mathcal{O} \rightarrow \mathcal{C}_s(\bar{X}) \times \mathcal{C}_s(\bar{X}) \times \mathcal{O}^2 \rightarrow \mathcal{C}_s(\bar{X})^2 \times \mathcal{C}_s(\bar{X})^2 \rightarrow \mathcal{C}_s(\bar{X})^2$$

for some Zariski open neighborhood \mathcal{O} of a k -rational point $0 \in \mathbf{P}^1$, where $\sigma(Z_1, Z_2, Z_3, Z_4) = (Z_1 + Z_4, Z_2 + Z_3)$.

We obtain a rational equivalence relating Z' and Z'' through cycles each of which meet Y properly as follows. The equivalence is parametrized by

$$R \equiv (\{(1, 0)\} \times \mathcal{O}) \cup (T \times \{p\}) \cup (\{(0, 1)\} \times \mathcal{O}).$$

On $\{(1, 0)\} \times \mathcal{O}$ (respectively, $\{(0, 1)\} \times \mathcal{O}$), we consider the restriction of Θ and then throw away those irreducible components whose fibres above $\bar{X} \times \{(1, 0)\} \times \{0\}$ (resp., $\bar{X} \times \{(0, 1)\} \times \{0\}$) do not meet Y properly. The result gives well defined rational equivalences starting from Z' and Z'' in view of the additivity of Θ . On $T \times \{p\}$ we simply take the

restriction of Θ . Thus, intersecting this rational equivalence with Y determines a rational equivalence between $Y \bullet Z', Y \bullet Z''$ parametrized by R .

Suppose now that k is an arbitrary finite field and $k \rightarrow k'$ is a finite field extension with k' sufficiently large. Let $\pi : X_{k'} \rightarrow X$ denote the finite, flat map given by base extension, where $X_{k'} \equiv X \times_{\text{Spec}(k)} \text{Spec}(k')$; this is a map of algebraic k schemes, but not a map of quasi-projective algebraic varieties over k . By the preceding argument, the rational equivalence class of $\pi^*(Y \bullet Z' - Y \bullet Z'')$ is 0 in the Chow group of $r + s - m$ -cycles on $X_{k'}$. Since $\pi_* \circ \pi^*$ equals multiplication by $d' = [k' : k]$ on cycles, $\pi_* \circ \pi^*$ also equals multiplication by d' on rational equivalence classes of cycles. By considering $\pi^*(Y \bullet Z' - Y \bullet Z'')$ for two such sufficiently large field extensions $k \rightarrow k', k \rightarrow k''$ of relatively prime degrees d', d'' over k , we conclude that $Y \bullet Z', Y \bullet Z''$ are rationally equivalent on X . \square

In [F-G], an intersection product was defined on the Lawson homology groups of a smooth, complex, quasi-projective variety X of dimension m :

$$L_r H_{2r+i}(X) \otimes L_s H_{2r+j}(X) \longrightarrow L_{r+s-m} H_{2(r+s-m)+i+j}(X), \quad r + s \geq m.$$

This product was defined using an argument involving the Fulton-MacPherson ‘‘Deformation to the normal cone’’ technique together with a homotopy lifting argument. The following consequence of Theorem 3.1 gives a more concrete construction of this intersection product for X projective as well as smooth. Even in this case, the construction of [F-G] remains useful in establishing the numerous good properties of this product.

Theorem 3.5 *Let X be a smooth, complex projective variety of dimension m and let r, s be non-negative integers with $r + s \geq m$. Let $\mathcal{Z}_r(X)$ denote the topological abelian group of r -cycles on X obtained as the ‘‘naive’’ group completion of the abelian monoid $\mathcal{C}_r(X)$ provided with the analytic topology. Then the pairing on homotopy groups induced by the intersection product of [F-G]*

$$\pi_i(\mathcal{Z}_r(X)) \otimes \pi_j(\mathcal{Z}_s(X)) \xrightarrow{\bullet} \pi_{i+j}(\mathcal{Z}_{r+s-m}(X))$$

can be represented as follows: given homotopy classes $\alpha \in \pi_i(\mathcal{Z}_r(X)), \beta \in \pi_j(\mathcal{Z}_s(X))$, there exist representative (base point preserving) maps $a : S^i \rightarrow \mathcal{Z}_r(X), b : S^j \rightarrow \mathcal{Z}_s(X)$ such that $a(t), b(u)$ intersect properly for all $t \in S^i, u \in S^j$ and such that the map sending $t \wedge u$ to the intersection of $a(t)$ and $b(u)$ is a continuous map $a \cdot b : S^{i+j} \rightarrow \mathcal{Z}_{r+s-m}(X)$ which represents $\alpha \bullet \beta$.

Proof. Since the topology on $\mathcal{Z}_k(X)$ is compactly generated by the images of $\mathcal{C}_{k, \leq e}(X)^2$ for $e > 0$, we may choose e sufficiently large that representatives a, b' of α, β lift to (not necessarily continuous) maps $\tilde{a} : \tilde{S}^i \rightarrow \mathcal{C}_{r, \leq e}(X)^2, \tilde{b}' : \tilde{S}^j \rightarrow \mathcal{C}_{s, \leq e}(X)^2$.

The continuous algebraic map Ψ of Theorem 3.1 chosen for this e determines a continuous map (for the analytic topology)

$$\Psi : \mathcal{Z}_s(X) \times \mathcal{O} \rightarrow \mathcal{Z}_s(X)$$

with the property that ψ_0 (i.e., the restriction of Ψ to $\mathcal{Z}_s(X) \times \{0\}$) is the identity. Define

$$b = \Psi \circ (b' \times i_p) : S^j \rightarrow \mathcal{Z}_s(X) \times \mathcal{O} \rightarrow \mathcal{Z}_s(X)$$

for some $p \in \mathcal{O} - \{0\}$, where $i_p : S^j \rightarrow \mathcal{O}$ is the constant map with value p . By Theorem 3.1.d and our choice of e , the r -cycles $a(t), b(u)$ intersect properly for all $t \in S^i, u \in S^j$.

The intersection of cycles meeting properly is continuous, so that a, b determine a continuous map $a \cdot b : S^{i+j} \rightarrow \mathcal{Z}_{r+s-m}(X)$. The fact that this map represents $\alpha \bullet \beta$ follows from [F-G;3.5.a]. \square

Remark 3.6. Our techniques fail to provide a representation of the intersection product on Lawson homology group of complex, smooth varieties X which are quasi-projective but not projective. This is because the “moves” we construct for s -cycles on a smooth completion \overline{X} of X do not restrict to moves on s -cycles on $\overline{X} - X$.

The following special case of Theorem 3.1 captures the fundamental assertion of the duality theory in [F-L2].

Theorem 3.7 *Let X be a smooth projective variety, $f : X \rightarrow B$ a flat morphism of varieties over k , and s an integer $\geq \dim(X) - \dim(B)$. Assume that k is an infinite field. Then for any positive integer e , there exist a Zariski open neighborhood $\mathcal{O} \subset \mathbf{P}^1$ of a k -rational point $0 \in \mathbf{P}^1$ and continuous algebraic maps*

$$\Psi = (\Psi^+, \Psi^-) : \mathcal{C}_s(X) \times \mathcal{O} \rightarrow \mathcal{C}_s(X)^2$$

satisfying the following properties (where ψ_p^\pm denotes the restriction of Ψ^\pm to $\mathcal{C}_s(X) \times \{p\}$):

- (a) $Z = \psi_0^+(Z) - \psi_0^-(Z)$ for every effective s -cycle Z on X .
- (b) For any effective s -cycle Z , the restriction of Ψ to $\{Z\} \times \mathcal{O}$ determines a rational equivalence $Z_{\mathcal{O}} \subset X \times \mathbf{P}^1$ whose fibre above $p \in \mathcal{O}$ equals $\psi_p^+(Z) - \psi_p^-(Z)$.
- (c) For every effective s -cycle Z of degree $\leq e$, the cycles $\psi_p^+(Z), \psi_p^-(Z)$ intersect properly each fibre $f^{-1}(b), b \in B$ of f , every $p \in \mathcal{O} - \{0\}$.

Proof. Apply Theorem 3.1 with e replaced by the larger of the integer e of the statement of this theorem and the maximum of the degrees of the fibres $f^{-1}(b), b \in B$ of f . \square

As remarked by Andrei Suslin, one can replace \mathcal{O} in Theorem 3.1 by \mathbf{A}^1 with only a bit more effort. We conclude by sketching the argument for the following “improvement” of Theorem 3.1.

Theorem 3.8 *Retain the hypotheses and notation of Theorem 3.1. Then there exists a continuous algebraic map*

$$\tilde{\Psi} = (\tilde{\Psi}^+, \tilde{\Psi}^-) : \mathcal{C}_s(X) \times \mathbf{A}^1 \rightarrow \mathcal{C}_s(X)^2$$

satisfying the following properties (where $\tilde{\psi}_p^\pm = \tilde{\Psi}^\pm|_{\mathcal{C}_s(X) \times \{p\}}$):

- (\tilde{a}) For every effective s -cycle Z ,

$$Z = \tilde{\psi}_0^+(Z) - \tilde{\psi}_0^-(Z).$$

- (\tilde{b}) For any k -rational point $p \in \mathbf{A}^1$, $\tilde{\psi}_p^\pm$ determines a pair of linear maps on effective s -cycles.
- (\tilde{c}) For any effective s -cycle Z on X , the restriction of $\tilde{\Psi}$ to $\{Z\} \times \mathbf{A}^1$ determines a rational equivalence $Z_\psi \subset \mathbf{P}^n \times \mathbf{P}^1$ whose fibre above a point $p \in \mathbf{A}^1$ is $\psi_p^+(Z) - \psi_p^-(Z)$.
- (\tilde{d}) For all effective cycles Y, Z on X of dimensions r, s and degrees $\leq e$ and all k -rational points $p \in \mathbf{A}^1 - \{0\}$, any component of excess dimension (i.e., $> r + s - m$) of either $|Y| \cap |\tilde{\psi}_p^+(Z)|$ or $|Y| \cap |\tilde{\psi}_p^-(Z)|$ lies in the singular locus of X .

Proof. We begin with $\Psi = (\Psi^+, \Psi^-)$ obtained in Theorem 3.1. Let $g : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be any morphism such that $g^{-1}(\infty) = \mathbf{P}^1 - \mathcal{O}$ and such that 0 occurs with multiplicity 1 in $g^{-1}(0)$; in particular, g restricts to a finite map $g : \mathcal{O} \rightarrow \mathbf{A}^1$. For any $Z \in \mathcal{C}_s(X)$ and $p \in \mathbf{A}^1$, define

$$\begin{aligned}\tilde{\Psi}_p^+(Z) &= \sum_{t \in g^{-1}(p)} \psi_t^+(Z) + \sum_{0 \neq t' \in g^{-1}(0)} \psi_{t'}^-(Z) \\ \tilde{\Psi}_p^-(Z) &= \sum_{t \in g^{-1}(p)} \psi_t^-(Z) + \sum_{0 \neq t' \in g^{-1}(0)} \psi_{t'}^+(Z).\end{aligned}$$

Using techniques as in [F], one readily checks that this is a continuous algebraic map. Namely, one first defines a rationally defined map by this formula applied to generic Z 's (i.e., associated to generic points of irreducible components of $\mathcal{C}_s(X)$). The closure of the graph of this rational map gives a continuous algebraic map since the value at any non-generic Z is given by the above formula and is thus independent of "path of specialization". The definition of $\tilde{\Psi}$ and properties (a)–(d) for Ψ proved in Theorem 3.1 immediately imply properties (\tilde{a})–(\tilde{d}) for $\tilde{\Psi}$. \square

References

- [Chev] C. Chevalley, Anneaux de Chow et Applications, Séminaire Chevalley, Secrétariat Math., Paris (1958).
- [Chow] W.S. Chow, On the equivalence classes of cycles in an algebraic variety, *Annals of Math.* **64** (1956), 450-479.
- [F] E. Friedlander, Algebraic cocycles, Chow varieties, and Lawson homology, *Compositio Math.* **77** (1991), 55-93.
- [F-G] E. Friedlander and O. Gabber, Cycles spaces and intersection theory, in *Topological Methods in Modern Mathematics* (1993), 325-370.
- [F-L] E. Friedlander and H.B. Lawson, A theory of algebraic cocycles, *Annals of Math.* **136** (1992), 361-428.
- [F-L2] E. Friedlander and H.B. Lawson, Duality relating cycle and cocycle spaces, *Topology* **36** (1997), 533-565.
- [F-V] E. Friedlander and V. Voevodsky. Bivariant cycle cohomology. To appear in *Annals of Math Studies*.
- [Fu] W. Fulton, *Intersection Theory*, Springer-Verlag, New York, 1984.

- [L] H.B. Lawson, Algebraic cycles and homotopy theory, *Annals of Math.* **129** (1989), 253-291.
- [L-F] P. Lima-Filho, Completions and fibrations for topological monoids and excision for Lawson homology, *Trans. A.M.S.* **340** (1993), no. 1, 127-147.
- [Mat] H. Matsumura, *Commutative Algebra*, Benjamin, 1970.
- [Mu] D. Mumford, *The Red Book of Varieties and Schemes*, Lecture Notes in Math 1358, Springer-Verlag, New York 1988.
- [R] J. Roberts, Chow's Moving Lemma, in *Algebraic Geometry Oslo 1970* (1972), 89-96.
- [S1] P. Samuel, *Méthodes d'algèbre abstraite en géométrie algébrique*, Ergebnisse der math., Springer-Verlag, 1955.
- [S2] P. Samuel, Rational equivalence of arbitrary cycles, *Amer. J. Math.* **78** (1956), 383-400.
- [S3] P. Samuel, Séminaire sur l'équivalence rationnelle, Orsay Seminar, 1971.