

Error Analysis of a Computation of Euler's Constant*

By W. A. Beyer and M. S. Waterman

Abstract. A complete error analysis of a computation of γ , Euler's constant, is given. The results have been used to compute γ to 7114 places and this value has been deposited in the UMT file.

1. **Introduction.** In a paper on ergodic computations with continued fractions [1], we used 3561 decimal places of γ , Euler's constant, as given by Sweeney [7] to compute 3420 partial quotients of the continued fraction expansion of γ . The partial quotients were sent to the Unpublished Manuscript Tables file and were there compared by Dr. Wrench with those given by Choong et al. [3]. Some disagreements were found and it was eventually decided to recompute Sweeney's value. This involved a careful reading of Sweeney's method and, as his error analysis is not detailed, a distinct error analysis resulted. This analysis is presented here.

2. **Error Analysis.** We begin with the exponential integral $-\text{Ei}(-x)$ [2, p. 334], and we consider only $x > 1$:

$$(1) \quad -\text{Ei}(-x) = \int_x^\infty \frac{e^{-t}}{t} dt = -\gamma - \ln x + S(x),$$

where

$$S(x) = x - \frac{x^2}{2 \cdot 2!} + \frac{x^3}{3 \cdot 3!} - \frac{x^4}{4 \cdot 4!} + \dots$$

The analysis of [6, p. 26] can be adapted to show

$$\int_x^\infty \frac{e^{-t}}{t} dt = \frac{e^{-x}}{x} \left(1 - \frac{1!}{x} + \frac{2!}{x^2} - \dots + \frac{(-1)^n n!}{x^n} + R_n(x) \right),$$

where $|R_n(x)| \leq (n+1)!/x^{n+1}$. However, we only require $n = 0$ and it is easy to see that

$$xe^x \int_x^\infty \frac{e^{-t}}{t} dt = \int_0^\infty \frac{e^{-s}}{1+s/x} ds = 1 - \int_0^\infty \frac{s/x}{1+s/x} e^{-s} ds = 1 + R_0(x).$$

Since, for $x > 0$,

Received June 26, 1973.

AMS (MOS) subject classifications (1970). Primary 10-04, 10A40; Secondary 41-04, 41A25.

* Work partly performed under the auspices of the U. S. Atomic Energy Commission while one of the authors (M.S.W.) was a faculty participant of the Associated Western Universities at Los Alamos Scientific Laboratory. The work was also supported in part by NSF grants GP-28313 and GP-28313 #1.

Copyright © 1974, American Mathematical Society

$$0 < \int_0^{\infty} \frac{s/x}{1+s/x} e^{-s} ds < \frac{1}{x} \int_0^{\infty} s e^{-s} ds = \frac{1}{x},$$

we infer that

$$(2) \quad \frac{e^{-x}}{x} - \frac{e^{-x}}{x^2} \leq \int_x^{\infty} \frac{e^{-t}}{t} dt \leq \frac{e^{-x}}{x}.$$

By Eqs. (1) and (2),

$$(3) \quad S(x) - \frac{e^{-x}}{x} - \ln x \leq \gamma \leq S(x) + \frac{e^{-x}}{x^2} - \frac{e^{-x}}{x} - \ln x.$$

Our problem is to use Eq. (3) to compute γ to a desired number of decimal places. After x is taken to be a power of 2, we must approximate e^{-x}/x , $\ln 2$, and $S(x)$. The computation was done on the Maniac II computer which does multiple-precision integer arithmetic without special programming. Therefore each function above will be multiplied by an appropriate power of 10, say 10^α . Of the α places in our answer, we will require that each answer be correct to $d - 1$ places. Equation (3) becomes

$$(4) \quad 10^\alpha S(x) - 10^\alpha \frac{e^{-x}}{x} - 10^\alpha \ln x \leq 10^\alpha \gamma \leq 10^\alpha S(x) + 10^\alpha \frac{e^{-x}}{x^2} - 10^\alpha \frac{e^{-x}}{x} - 10^\alpha \ln x.$$

We first consider the error in the exponential terms of (4).

$$\left| 10^\alpha \left(\frac{e^{-x}}{x} - \frac{e^{-x}}{x^2} \right) \right| \leq 10^\alpha \frac{e^{-x}}{x} < 10^{\alpha - x/\ln 10}.$$

If the exponential terms are neglected in (3) and we desire $d - 1$ correct places, we must have $\alpha - x/\ln 10 < \alpha - d$ or $d \ln 10 < x$. Thus, we determine d from

$$(5) \quad d = \lceil x/\ln 10 \rceil.$$

The following procedure is used to approximate $S(x)$. Let

$$A_{n-1} = 10^\alpha - \frac{n-1}{n^2} x,$$

$$A_k = 10^\alpha - \frac{k}{(k+1)^2} x A_{k+1}, \quad 1 \leq k < n-1.$$

Then define $T(x)$ by

$$10^\alpha T(x) = x A_1 = x \left(10^\alpha - \frac{x}{2^2} A_2 \right)$$

$$= x \left(10^\alpha - \frac{x}{2^2} \left(10^\alpha - \frac{2x}{3^2} A_3 \right) \right)$$

$$\dots$$

$$= 10^\alpha \left(x - \frac{x^2}{2 \cdot 2!} + \frac{x^3}{3 \cdot 3!} - \frac{x^4}{4 \cdot 4!} + \dots + (-1)^{n+1} \frac{x^n}{n \cdot n!} \right).$$

The truncation error in using $10^\alpha T(x)$ in place of $10^\alpha S(x)$ is

$$\begin{aligned} |10^\alpha S(x) - 10^\alpha T(x)| &\leq 10^\alpha \left(\frac{x^{n+1}}{(n+1)(n+1)!} + \frac{x^{n+2}}{(n+2)(n+2)!} + \frac{x^{n+3}}{(n+3)(n+3)!} + \dots \right) \\ &\leq \frac{10^\alpha}{n+1} \left(\frac{x^{n+1}}{(n+1)!} + \frac{x^{n+2}}{(n+2)!} + \frac{x^{n+3}}{(n+3)!} + \dots \right). \end{aligned}$$

The quantity in parentheses is the remainder term in the Taylor expansion of e^x and is therefore equal to $x^{n+1}e^{\theta x}/(n+1)!$, $\theta \in (0, 1)$. Next, we assume $n > 2x$ and use a technique of Courant [4, p. 326] to obtain $x^{n+1}/(n+1)! < (2x)^{2x}/(2x)!2^{-n-1}$. Thus

$$|10^\alpha S(x) - 10^\alpha T(x)| < \frac{10^\alpha}{n+1} e^x \left(\frac{(2x)^{2x}}{(2x)!} 2^{-n-1} \right).$$

Using the fact that Stirling's formula underestimates $(2x)!$ [5, p. 54], we obtain

$$|10^\alpha S(x) - 10^\alpha T(x)| < \frac{10^\alpha}{n+1} \frac{e^{3x-(n+1)\ln 2}}{(2\pi)^{1/2}(2x)^{1/2}} < 10^{\alpha+(3x-(n+1)\ln 2)/\ln 10}.$$

Since we require $d - 1$ correct places, we take

$$\alpha + (3x - (n + 1) \ln 2) / \ln 10 < \alpha - d$$

which yields

$$n > d \ln 10 / \ln 2 + 3x / \ln 2 - 1.$$

But we have $x > d \ln 10$, so it is sufficient to take

$$(6) \quad n = [4x / \ln 2].$$

We note that $n = [4x / \ln 2] > 2x$ as required above.

There is also round-off error in computing $10^\alpha T(x)$. Assume that an error of ϵ_k is made in the k th iteration:

$$A_k = 10^\alpha - \frac{kx}{(k+1)^2} A_{k+1} + \epsilon_k.$$

Then

$$\begin{aligned} 10^\alpha \hat{T}(x) &= x A_1 \\ &\dots \\ &= 10^\alpha T(x) + \left(x\epsilon_1 - \frac{x^2}{2 \cdot 2!} \epsilon_2 + \frac{x^3}{3 \cdot 3!} \epsilon_3 - \dots + \frac{(-1)^{n+1} x^n}{n \cdot n!} \epsilon_n \right). \end{aligned}$$

If we assume $|\epsilon_k| \leq \epsilon = 1$ for all k , then

$$|10^\alpha \hat{T}(x) - 10^\alpha T(x)| < e^x.$$

Now $e^x = 10^{x/\ln 10} < 10^{\alpha-d}$ if

$$(7) \quad \alpha = 2d + 1.$$

If γ to disc

decimal places, one can use (3) to compute. The computation of the decimals of $\ln 2$ is

mine

to be some positive integer to be determined (can be obtained from Taylor's series):

$$\left(\frac{1}{3} + \frac{10^\beta}{5 \cdot 3^5} + \frac{10^\beta}{7 \cdot 3^7} + \dots \right).$$

This

$$\left[\dots + \left[\frac{10^\beta}{(2(k-1)+1)3^{2(k-1)+1}} \right] \right]$$

where

defined β by the condition that

(8)

$$(2k+1)3^{2k+1}.$$

The

(9)

$$\left(\left[\frac{10^\beta}{3 \cdot 3^3} \right] \right) \left(\frac{1}{(2k+1)3^{2(k-1)+1}} - \left[\frac{10^\beta}{(2(k-1)+1)3^{2(k-1)+1}} \right] \right) \left(\frac{1}{(2k+3)3^{2k+3}} + \dots \right).$$

The

(9) is dominated by

$$\frac{1}{(2k+1)3^{2k+1}} \frac{1}{8} \leq \frac{9}{4},$$

max

(9) are dominated by $k - 1$. Hence

(10)

$$k \leq 2(k-1) + 9/4,$$

where by

(10) satisfies (8). An upper bound to k as given

(11)

$$\frac{\ln 10}{\ln 3}.$$

He

(12)

$$k \leq \beta \ln 10 / \ln 3 + \frac{1}{4}.$$

In as and

(12). One sees from (12) that the error in $\ln 2$ is 10^{-k} places. We actually have only reported

TABLE 1

n	Sample Frequency of n	Theoretical Frequency of n :
		$\frac{1}{\ln 2} \ln \frac{(n+1)^2}{n(n+2)}$
1	0.4225	0.4150
2	0.1646	0.1699
3	0.0896	0.0931
4	0.0527	0.0589
5	0.0438	0.0406
6	0.0308	0.0297
7	0.0228	0.0227
8	0.0216	0.0179
9	0.0121	0.0144
10	0.0124	0.0119

TABLE 2

x	Guaranteed Number of Correct Digits $d - 1$	Actual Number of Correct Digits
8	2	4
16	5	7
32	12	14
64	26	29
128	54	57
256	110	113
512	221	224
1024	443	446
2048	888	889
4096	1777	1795
8192	3556	3561
16384	7114	—

4. Computation of γ . For our calculation, we used $x = 2^{14}$. From this x , we obtained $d = [x/\ln 10] = 7115$, $\alpha = 2d + 1 = 14231$, $n = [4x/\ln 2] = 94548$, and $k = [\alpha \ln 10 / (2 \ln 3)] + 1 = 14914$. The above analysis shows that our computation of γ is accurate to 7114 places. The errors from e^{-x}/x and $S(x)$ might each affect the 7115th place.

From this computation, we obtained 7114 correct decimal places of γ . These values were used to calculate 6920 partial quotients in the continued fraction expansion of γ . The 7121 places of $\ln 2$ yielded 6890 partial quotients of $\ln 2$. Note that

the number of partial quotients of γ is more than that of $\ln 2$. These have been sent to the Unpublished Manuscript Tables (UMT) file of this journal.

In Choong et al. [3], Table 1 gives sample frequency of n and theoretical frequency of n for 3470 partial quotients of γ . Our Table 1 corrects their Table 1. Our Table 2 gives our results for $x = 2^t$ ($t = 3, 4, \dots, 14$) and is thus a check of our analysis.

Acknowledgements. The authors thank Dr. J. W. Wrench, Jr. for helpful suggestions and encouragement in this work. We thank the referee for pointing out some oversights.

Los Alamos Scientific Laboratory
Los Alamos, New Mexico 87544

Idaho State University
Pocatello, Idaho 83201

1. W. A. BEYER & M. S. WATERMAN, "Ergodic computations with continued fractions and Jacobi's algorithm," *Numer. Math.*, v. 19, 1972, pp. 195-205. Errata, v. 20, 1973, p. 430. MR 46 #2832.
2. T. J. F. A. BROMWICH, *An Introduction to the Theory of Infinite Series*, 2nd rev. ed., Macmillan, London, 1949.
3. K. Y. CHOONG, D. E. DAYKIN & C. R. RATHBONE, "Rational approximations to π ," *Math. Comp.*, v. 25, 1971, pp. 387-392. (See also *Math. Comp.*, v. 27, 1973, p. 451, MTE 501.) MR 46 #141.
4. R. COURANT, *Differential and Integral Calculus*. Vol. I, 2nd rev. ed., Interscience, New York, 1937.
5. W. FELLER, *An Introduction to Probability Theory and its Applications*. Vol. I, 3rd rev. ed., Wiley, New York, 1968. MR 37 #3604.
6. G. H. HARDY, *Divergent Series*, Clarendon Press, Oxford, 1949. MR 11, 25.
7. D. W. SWEENEY, "On the computation of Euler's constant," *Math. Comp.*, v. 17, 1963, pp. 170-178. (See Corrigendum in *Math. Comp.*, v. 17, 1963, p. 488.) MR 28 #3522.