## **MATH 445**

Computer Project 1 Due Friday, October 21, 2022

You are free to use any programming language or environment and any help you want.

Please, use at least a 10pt font, and do not submit more than 10 pages of printouts for this assignment. When you are done, please print everything into a single PDF file and upload the file to Blackboard.

**Problem 1.** The main objective is to practice plotting graphs. PART 1. Plot the graph of the function

$$h_{10}(x) = \sum_{k=1}^{10} \frac{\sin\left((k!)^2 x\right)}{k!}$$

for  $x \in [0, 1], x \in [0, 0.1], x \in [0, 0.01], x \in [0, 0.001].$ 

What you will turn in:

- (1) Four separate graphs.
- (2) Printout of the program you used to generate the graphs.

Each page you turn in must have your name and date *printed* on it. Each graph must have a title, labeled axes, and the scale along each axis.

PART 2. Consider the function

$$h(x) = \sum_{k=1}^{\infty} \frac{\sin\left((k!)^2 x\right)}{k!}.$$

For what  $x \in [0, 1]$  will this function be

- (1) defined?
- (2) continuous?
- (3) differentiable?

Provide a one-sentence explanation for each of your answers.

**Problem 2.** The objective is to compute Fourier coefficients numerically and to analyze the Gibbs phenomenon. Let f = f(x) be a  $2\pi$ -periodic function defined for  $x \in (-\pi, \pi]$  by f(x) = x. Let  $S_n(x) = \sum_{k=1}^n b_k \sin(kx)$ , where  $b_k = (1/\pi) \int_{-\pi}^{\pi} f(x) \sin(kx) dx$ .

Do the following:

- (1) Plot the graphs of  $S_n(x)$  for n = 10, 50, 100 and  $x \in [-\pi, \pi]$ . The choice of the procedure to compute  $b_k$  is up to you. Keep in mind that if you divide the interval  $[-\pi, \pi]$  to approximate the integral, your step size must be small enough to "see" the oscillations of the sines.
- (2) Estimate  $\max_{x \in [-\pi,\pi]} S_n(x)$  for n = 10, 50, 100.
- (3) Compute  $\lim_{n\to\infty} \frac{S_n(\pi-\pi/n)}{\pi}$  (either compute analytically or guess from the graphs). This is a quantitative measure of the Gibbs phenomenon.

What you will turn in:

- (1) Three separate graphs with the corresponding values of  $\max_{x \in [-\pi,\pi]} S_n(x)$ .
- (2) Printout of the program you used to generate the graphs. Please indicate the procedure you used to compute the Fourier coefficients  $b_k$ .
- (3) The numerical value of  $\lim_{n\to\infty} \frac{S_n(\pi-\pi/n)}{\pi}$  and the corresponding explanations.

Each page you turn in must have your name and date *printed* on it. Each graph must have a title, labeled axes, and the scale along each axis.

Instructor: Sergey Lototsky, KAP 248D.

## Computer Project 2 Due Friday, December 2, 2022

You are free to use any programming language or environment and any help you want.

Please, use at least a 10pt font, and do not submit more than 12 pages of printouts for this assignment. When you are done, please print everything into a single PDF file and upload the file to Blackboard.

The objective of this assignment is to see how implicit numerical schemes work for parabolic and hyperbolic equations.

**Problem 1.** Consider the heat equation

$$u_t(x,t) = 0.25u_{xx}(x,t), \ 0 < t \le 2, \ 0 < x < 1,$$

with u(0,t) = u(1,t) = 0 and

$$u(x,0) = \begin{cases} 20x, & 0 \le x \le 1/2\\ 20(1-x), & 1/2 \le x \le 1. \end{cases}$$

Solve it numerically by the Crank-Nicholson method taking h = k = 0.1. Plot a 3-D graph of the result. Then compare the result with the Fourier series solution (use your judgement as to how many terms to keep in the Fourier series: this is your only chance to get to the exact solution as close as possible).

What you will turn in:

- (1) The graph  $(x, t, \bar{u}(x, t))$ , where  $\bar{u}$  is the numerical solution you got.
- (2) The graph  $(x, t, |\bar{u}(x, t) u(t, x)|)$ , where u is the Fourier series solution.
- (3) Printout of the program you used to generate the graphs.

Each page you turn in must have your name and date *printed* on it. The graph must have a title, labeled axes, and the scale along each axis.

Problem 2. Consider the wave equation

$$u_{tt}(x,t) = u_{xx}(x,t), \ 0 < t \le 2, \ 0 < x < 1,$$

with  $u(0,t) = u(1,t) = u_t(x,0) = 0$ , u(x,0) = x(1-x).

Solve it numerically by the implicit method taking h = k = 0.1. Plot a 3-D graph of the result. Then compare the result with the exact solution (unlike the heat equation, you can get the exact solution without the Fourier series.) What you will turn in:

- (1) The graph  $(x, t, \bar{u}(x, t))$ , where  $\bar{u}$  is the numerical solution you got.
- (2) The graph  $(x, t, |\bar{u}(x, t) u(t, x)|)$ , where u is the exact solution. This time, you have at least two choices for the exact solution: you can either truncate the Fourier series of the solution or you can use d'Alembert's formula and get the truly exact solution in the form

$$u(t,x) = \frac{S_{f,s}(x+ct) + S_{f,s}(x-ct)}{2}$$

where  $S_{f,s}$  is the Fourier sine series of the function f(x) = x(1-x).

(3) Printout of the program you used to generate the graphs.

Each page you turn in must have your name and date *printed* on it. The graph must have a title, labeled axes, and the scale along each axis.