

# Convergence of Random Variables<sup>1</sup>

**The setting:** probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , random variables  $\xi_n = \xi_n(\omega)$ ,  $n \geq 1$ , and  $\xi = \xi(\omega)$ ,  $\omega \in \Omega$ .

## Definitions.

- (1) **Uniform convergence:**  $\xi_n \rightrightarrows \xi$  if  $\forall \varepsilon > 0 \exists N = N(\varepsilon) \forall n > N \forall \omega \in \Omega : |\xi_n(\omega) - \xi(\omega)| < \varepsilon$ .
- (2) **Point-wise convergence:**  $\xi_n \xrightarrow{p.w.} \xi$  if  $\xi_n(\omega) \rightarrow \xi(\omega)$  for all  $\omega \in \Omega$ , that is,  
 $\forall \varepsilon > 0 \forall \omega \in \Omega \exists N = N(\varepsilon, \omega) \forall n > N : |\xi_n(\omega) - \xi(\omega)| < \varepsilon$ .
- (3) **Complete convergence:**  $\xi_n \xrightarrow{c.c.} \xi$  if  $\sum_{n=1}^{\infty} \mathbb{P}(|\xi_n - \xi| > \varepsilon) < +\infty$  for every  $\varepsilon > 0$ .
- (4) **Almost sure convergence:**  $\xi_n \xrightarrow{a.s.} \xi$  if  $\mathbb{P}\left(\omega : \lim_{n \rightarrow \infty} \xi_n(\omega) = \xi(\omega)\right) = 1$ . Also known as convergence with probability one.
- (5) **Convergence in  $L_p$ ,  $p > 0$ :**  $\xi_n \xrightarrow{L_p} \xi$  if  $\lim_{n \rightarrow \infty} \mathbb{E}|\xi_n - \xi|^p = 0$ .
- (6) **Convergence in probability:**  $\xi_n \xrightarrow{\mathbb{P}} \xi$  if  $\lim_{n \rightarrow \infty} \mathbb{P}(|\xi_n - \xi| > \varepsilon) = 0$  for every  $\varepsilon > 0$ .
- (7) **Convergence in distribution:**  $\xi_n \xrightarrow{d} \xi$  if  $\lim_{n \rightarrow \infty} \varphi_{\xi_n}(t) = \varphi_{\xi}(t)$  for every  $t \in \mathbb{R}$ ;  $\varphi_{\xi}(t) = \mathbb{E}e^{it\xi}$  is the characteristic function of  $\xi$ .
- (8) **Uniform integrability:** the sequence  $\{\xi_n, n \geq 1\}$  is uniformly integrable (UI) if

$$\lim_{a \rightarrow +\infty} \sup_{n \geq 1} \mathbb{E}\left(|\xi_n| 1(|\xi_n| > a)\right) = 0.$$

Each of the following is a sufficient condition for UI:

- $|\xi_n| \leq \eta$  for all  $n$  and  $\mathbb{E}\eta < +\infty$ ;
- $\mathbb{E} \sup_n |\xi_n| < \infty$ ;
- $\sup_n \mathbb{E}g(|\xi_n|) < \infty$  for a non-negative increasing function  $g = g(x)$ ,  $x \geq 0$ ,  
such that  $\lim_{x \rightarrow +\infty} \frac{g(x)}{x} = +\infty$ . For example,  $\sup_n \mathbb{E}|\xi_n|^r < \infty$  for some  $r > 1$ .

## General implications:

$$\xrightarrow{p.v.} \Leftarrow \Leftrightarrow \Rightarrow \xrightarrow{c.c.} \Rightarrow \xrightarrow{a.s.} \Rightarrow \xrightarrow{\mathbb{P}} \Rightarrow \xrightarrow{d}; \quad \xrightarrow{L_p} \Rightarrow \xrightarrow{\mathbb{P}}.$$

**The (non-trivial) reasons.**  $\Rightarrow \Rightarrow \xrightarrow{c.c.}$  because  $\Rightarrow$  implies  $\mathbb{P}(|\xi_n - \xi| > \varepsilon) = 0$  for all sufficiently large  $n$ ;  
 $\xrightarrow{c.c.} \Rightarrow \xrightarrow{a.s.}$  by the first Borel-Cantelli;  $\xrightarrow{a.s.} \Rightarrow \xrightarrow{\mathbb{P}}$  because  $\xrightarrow{a.s.}$  means  $\lim_{n \rightarrow \infty} \mathbb{P}(\sup_{k \geq n} |\xi_k - \xi| > \varepsilon) = 0$ ;  
 $\xrightarrow{\mathbb{P}} \Rightarrow \xrightarrow{d}$  by the Dominate Convergence Theorem;  $\xrightarrow{L_p} \Rightarrow \xrightarrow{\mathbb{P}}$  by Markov's inequality.

**Basic counterexamples**, with  $\Omega = [0, 1]$  and  $\mathbb{P}$  the Lebesgue measure,

- (1) **the Growing Pulse**  $\xi_n(x) = n^\alpha 1(0 < x < n^{-\beta})$  or  $\xi_n(x) = n^\alpha 1(0 \leq x < n^{-\beta})$  [if DO NOT want  $\xi_n$  to converge to zero point-wise], for suitable  $\alpha > 0, \beta > 0$ , shows that
  - $\xrightarrow{p.v.}$  or  $\xrightarrow{a.s.} \not\Rightarrow \xrightarrow{L_p}$  or  $\xrightarrow{c.c.}$
  - $\xrightarrow{c.c.} \not\Rightarrow \Rightarrow$  or  $\xrightarrow{p.w.}$
- (2) **the Typewriter Sequence**  $\xi_{n,k} = 1(k2^{-n} < x < (k+1)2^{-n})$ ,  $k = 0, \dots, 2^n - 1$ ,  $n = 0, 1, 2, \dots$ , shows that

$$\xrightarrow{L_p}, \quad 0 < p < \infty \not\Rightarrow \xrightarrow{a.s.}$$

## Special implications

- (1)  $\xrightarrow{\mathbb{P}} \Rightarrow \xrightarrow{a.s.}$  if the sequence  $\{\xi_n, n \geq 1\}$  is monotone;
- (2)  $\xrightarrow{d} \Rightarrow \xrightarrow{\mathbb{P}}$  if the limit  $\xi$  is non-random: there exists a real number  $c$  such that  $\mathbb{P}(\xi = c) = 1$ ;
- (3)  $\xrightarrow{d} \Rightarrow \xrightarrow{a.s.}$  for sums of independent random variables [using Cauchy criterion and zero-one law];

<sup>1</sup>Sergey Lototsky, USC

- (4)  $\xrightarrow{\mathbb{P}} \Rightarrow \xrightarrow{L_p}$  if the sequence  $\{|\xi_n|^p, n \geq 1\}$  is UI.
- (5)  $\xrightarrow{a.s.} \Rightarrow \xrightarrow{c.c.}$  if the random variables  $\xi_n$  are independent AND the limit  $\xi$  is non-random [by the second Borel-Cantelli; if the limit  $\xi$  is random, then the random variables  $\xi_n - \xi$  might be dependent and the second Borel-Cantelli will not apply];
- (6) If  $\xi_n$  are iid and  $\mathbb{E}|\xi_n| < \infty$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \xi_k = \mathbb{E}\xi_1$  both with probability one and in  $L_1$ .
- (7) If  $\xi_n \xrightarrow{\mathbb{P}} \xi$ , then  $\xi_{n_k} \xrightarrow{c.c.} \xi$  along a suitable sub-sequence;
- (8) If  $\xi_n \xrightarrow{\mathbb{P}} \xi$  and, for some  $p > 0$ ,  $\lim_{n \rightarrow \infty} \mathbb{E}|\xi_n|^p = \mathbb{E}|\xi|^p$ , then  $\{|\xi_n|^p, n \geq 1\}$  is UI and  $\xi_n \xrightarrow{L_p} \xi$ .

### Working with limits

- (1) If  $M = a.s.$  or  $M = \mathbb{P}$ , and  $\xi_n \xrightarrow{M} \xi$ ,  $\eta_n \xrightarrow{M} \eta$ , then  $\xi_n \pm \eta_n \xrightarrow{M} \xi \pm \eta$ ,  $\xi_n \eta_n \xrightarrow{M} \xi \eta$ , and, if  $\eta \neq 0$ , then also  $\xi_n/\eta_n \xrightarrow{M} \xi/\eta$ .
- (2) If  $\xi_n \xrightarrow{M} \xi$ , with  $M = a.s.$  or  $\mathbb{P}$  or  $d$ , and  $f = f(x)$ ,  $x \in \mathbb{R}$  is a continuous function, then  $f(\xi_n) \xrightarrow{M} f(\xi)$ . The result is obvious when  $M = a.s.$ , easy when  $M = \mathbb{P}$ , and non-trivial [known as the **Mann-Wald Theorem**] when  $M = d$ .
- (3) **Slutsky's Theorem:** if  $\xi_n \xrightarrow{d} \xi$  and  $\eta_n \xrightarrow{d} b \in \mathbb{R}$  non-random, then  $\xi_n \pm \eta_n \xrightarrow{d} \xi \pm b$ ,  $\xi_n \eta_n \xrightarrow{d} b\xi$ , and, if  $b \neq 0$ , then also  $\xi_n/\eta_n \xrightarrow{d} \xi/b$ .

### Further Facts

- $\xrightarrow{p.w.}$  corresponds to a topology, but the topology is not metrizable;
- $\xrightarrow{\mathbb{P}}$  is metrizable: the function  $\rho(\xi, \eta) = \mathbb{E} \frac{|\xi - \eta|}{1 + |\xi - \eta|}$  defines a metric on the space of random variables, and  $\xi_n \xrightarrow{\mathbb{P}} \xi$  if and only if  $\lim_{n \rightarrow \infty} \rho(\xi_n, \xi) = 0$ ;
- $\xrightarrow{d}$  is metrizable using the **Lévy-Prokhorov metric**;
- $\xrightarrow{a.s.}$  is not metrizable and does not correspond to any topology.